



DOI: 10.4274/jcrpe.galenos.2025.2025-6-23

J Clin Res Pediatr Endocrinol

Genetics of Idiopathic Hypogonadotropic Hypogonadism

✉ A. Kemal Topaloğlu¹, ✉ Leman Damla Kotan²

¹Harvard Medical School, Massachusetts General Hospital, Department of Pediatric Endocrinology, Boston, Massachusetts, USA

²Çukurova University Faculty of Medicine, Department of Pediatric Endocrinology, Adana, Türkiye

Cite this article as: Topaloğlu AK, Kotan LD. Genetics of idiopathic hypogonadotropic hypogonadism. J Clin Res Pediatr Endocrinol. [Epub Ahead of Print]

ABSTRACT

Idiopathic hypogonadotropic hypogonadism (IHH) comprises a group of disorders characterized by deficient secretion or action of gonadotropin-releasing hormone (GnRH), leading to impaired pubertal development and infertility. Traditionally, IHH is classified into Kallmann syndrome, associated with anosmia, and normosmic IHH, in which olfactory function is preserved. The condition exhibits marked genetic heterogeneity. Advances in next generation sequencing have significantly expanded the genetic landscape of IHH, with pathogenic variants identified in over 60 genes, accounting for up to 50% of cases. Oligogenic inheritance is increasingly recognized, occurring in 10-20% of individuals. The potential for spontaneous or treatment-induced clinical recovery in a subset of patients, along with phenotypic overlap with constitutional delay of growth and puberty, presents additional diagnostic challenges. Despite these complexities, genetic studies of IHH have provided critical insights into fundamental neuroendocrine processes, most notably the recent elucidation of the Kisspeptin, Neurokinin B, Dynorphin neurons as the GnRH pulse generator. These discoveries have also informed the development of targeted therapies, exemplified by the recent FDA approval of fezolinetant, a neurokinin B receptor antagonist, for the treatment of menopausal vasomotor symptoms.

Keywords: Hypogonadotropic hypogonadism, delayed puberty, genetics, etiology

Introduction

In vertebrates, gonadotropin releasing hormone (GnRH)-secreting neurons develop outside the central nervous system, originating from the nasal placode. They migrate along olfactory-derived vomeronasal axons to their final location in the hypothalamus (1).

The activity of the hypothalamic-pituitary-gonadal (HPG) axis demonstrates significant variability across the human lifespan (2). During early adolescence, a gradual reactivation of this neuroendocrine axis initiates the development of secondary sexual characteristics and the maturation of the

reproductive system, marking the onset of puberty. This complex developmental process typically starts around 10 to 11 years of age in girls and boys respectively and spans from two to five years. Epidemiological data suggest that approximately 50% to 75% of the variation in the age at onset of puberty is influenced by genetics (3). The failure to undergo pubertal progression results in sexual immaturity and infertility, a clinical state referred to as hypogonadism. When this condition arises from anatomical malformations or functional impairments that compromise the secretion of GnRH or the subsequent release of pituitary gonadotropins, it is specifically classified as hypogonadotropic hypogonadism (HH).

Corresponding Author: Prof. A. Kemal Topaloğlu, Harvard Medical School, Massachusetts General Hospital, Department of Pediatric Endocrinology, Boston, Massachusetts, USA

E-mail: atopaloglu@mgh.harvard.edu **ORCID:** orcid.org/0000-0003-2931-9552

Received: 24.06.2025 **Accepted:** 02.09.2025 **Epub:** 17.09.2025



Idiopathic Hypogonadotropic Hypogonadism

The term idiopathic HH (IHH) refers to a condition characterized by delayed or absent sexual maturation due to deficient secretion or action of GnRH, in the absence of any identifiable anatomical or functional cause. IHH is traditionally categorized into two main forms: Kallmann syndrome (KS), which is associated with anosmia or hyposmia, and normosmic IHH (nIHH), in which olfactory function remains intact. IHH may be either congenital or acquired, with congenital cases comprising the majority of those that have a hereditary basis. In female individuals, clinical signs typically do not become evident until the early adolescent years. In contrast, male infants may exhibit signs of reproductive dysfunction at birth due to the critical activity of the HPG axis between the sixteenth and twenty-second weeks of gestation, a period during which androgen production is essential for the proper virilization of the male fetus with a 46,XY karyotype. As a result, congenital IHH in males may present with micropenis and/or undescended testes (cryptorchidism) at birth. In some cases, the degree of under-virilization is sufficient to warrant evaluation for a “disorder of sexual development”. The brief reactivation of the HPG axis during early infancy, commonly referred to as “minipuberty”, occurring approximately between four and sixteen weeks of life, presents a valuable diagnostic window during which both male and female infants with congenital IHH may be identified (4). KS is typically attributed to aberrant embryonic development and/or disrupted migration of GnRH-secreting neurons. During embryogenesis, these GnRH neurons originate from the nasal placode and migrate along olfactory axons to reach the hypothalamus. Consequently, the close developmental association between GnRH and olfactory neurons underlies the characteristic clinical presentation of HH accompanied by anosmia or hyposmia. In addition to reproductive and olfactory deficits, individuals with KS frequently exhibit a spectrum of non-reproductive congenital anomalies, including cleft palate, unilateral renal agenesis, limb malformations such as split hand/foot malformation and shortened metacarpals, sensorineural hearing loss, and mirror movements (synkinesia) (5). In contrast nIHH refers to those IHH cases in which patients have an intact sense of smell (6). nIHH arises from dysfunction of the GnRH neurons that are properly located within the hypothalamus. These cases typically lack any associated congenital anomalies.

Caution is warranted when employing the classifications of KS and nIHH, as the distinction between these entities can be ambiguous. This is exemplified by mutations in the *FGFR1* gene, which could be associated with either phenotype. Mutations in *CCDC141* or *IGSF10*, although typically associated with nIHH, have been shown *in vitro* to impair the migration of GnRH neurons, an abnormality more commonly linked to the KS phenotype. These findings underscore the complexity of the underlying molecular

mechanisms and challenge the traditional dichotomy between KS and nIHH (7,8).

Pubertal delay is the most typical presentation of IHH. Pubertal delay is defined as absence of breast development (Tanner breast stage 1) in a girl at age 13 years or failure to achieve a testicular volume of 4 mL in a boy by age 14 years (9). By far the most common cause of delayed puberty is constitutional delay in growth and puberty (CDGP), also known as self-limited delayed puberty, which is not a disease *per se* but a maturational delay in development at the extreme of the population standards. CDGP accounts for pubertal delay in two third of boys and one third of girls (10). As CDGP is a diagnosis of exclusion, it must be carefully considered in the differential diagnosis of IHH. Differentiating between these two conditions frequently necessitates prolonged clinical observation and extensive diagnostic evaluation.

Studies have shown that some variants in established puberty-related genes, including *TAC3* and *TACR3*, are present in both individuals with nIHH and those with CDGP within the same families. These findings suggest that CDGP may represent a milder, transient manifestation of the same genetic defect underlying IHH, indicating a shared pathophysiological continuum between the two conditions (11). Clinicians often initiate a low-dose sex steroid regimen to “jump-start” pubertal development in patients suspected of having CDGP. It is now well established that approximately 10-20% of individuals with IHH experience clinical recovery, occurring either spontaneously or, more commonly, following sex steroid replacement therapy (12,13). These observations suggest that CDGP and IHH may share underlying pathophysiological mechanisms. This supports the concept of a phenotypic continuum ranging from normal pubertal timing to severe forms of IHH, with CDGP representing an intermediate point along this spectrum. On the other hand, a recent study found that the common genetic variants associated with pubertal timing in the general population contribute substantially to the genetic basis of CDGP, but only minimally to that of IHH (14). Furthermore, a more recent study involving 71 CDGP subjects revealed no mutations in genes associated with nIHH, such as *GNRHR*, *TAC3*, and *TAC3R*. This study revealed new candidate genes for CDGP, most notably *INHBB*, encoding the beta B subunit of inhibin, which is associated with age at menarche (15). In yet another study *MC3R* loss-of-function variants were overrepresented in patients with CDGP in comparison to IHH (16). Collectively, these recent studies suggest that the genetic architecture underlying CDGP and IHH may be distinct. Currently identified genetic defects explain up to 50% of all IHH cases (5,17). To date, mutations in nearly 60 genes have been implicated in the pathogenesis of IHH. A comprehensive list of currently known IHH-associated genes

is presented in Table 1. In a subset of patients or pedigrees, more than one pathogenic variant in different IHH-associated genes is identified, a phenomenon referred to as oligogenic inheritance or etiology. This mode of inheritance is estimated to account for 10-20% of all IHH cases (18,19,20,21). With the increased application of comprehensive, unbiased genetic approaches, such as whole exome sequencing (WES), it has become evident that oligogenic inheritance is more prevalent in Mendelian disorders than previously recognized (22). From the diagnostic point of view designing a panel of genes for targeted exome sequencing may prove to be practical in aiding timely differential diagnosis of delayed puberty. Such gene panels may prioritize genes more commonly implicated in patients with IHH or CDGP, and in our view, should at minimum include *FGFR1*, *ANOS1*, *CHD7*, *PROKR2*, *GNRHR*, *KISS1R*, *TAC3*, *TACR3*, *FGF8*, *FGF17*, *PROK2*, *CCDC141*, *SEMA3A*, *IGSF10*, *INHBB*, *MC3R*, and *IL17RD* (5,15,16,23,24).

Genes Associated with IHH

Kallmann syndrome (KS) associated genes

X-linked recessive, autosomal dominant (AD), and autosomal recessive (AR) inheritance patterns have all been described in association with KS. However, KS frequently occurs as a sporadic condition. Even in familial cases, considerable intrafamilial phenotypic variability is commonly observed, with individuals harboring the same genetic mutation exhibiting a wide range of clinical manifestation (25,26,27). Based on the presence of specific associated clinical features, genetic screening can be prioritized for particular gene(s): synkinesia (*ANOS1*), dental agenesis (*FGF8/FGFR1*), digital bony abnormalities (*FGF8/FGFR1*) and hearing loss (*CHD7*, *SOX10*) (28). A shared pathophysiological mechanism among genes implicated in KS involves the interaction of fibroblast growth factor signaling, prokineticin signaling, and Anosmin-1 with heparan sulfate glycosaminoglycan moieties within extracellular signaling complexes. These interactions are thought to facilitate the proper migration of GnRH neurons during embryonic development (29,30).

Table 1. The list of genes associated with idiopathic hypogonadotropic hypogonadism

Gene	HGNC ID	Approved name	OMIM phenotype	Phenotype MIM number
<i>AMH</i>	464	Anti-Müllerian hormone	Persistent Mullerian duct syndrome, type I	261550
<i>AMHR2</i>	465	Anti-Müllerian hormone receptor type 2	Persistent Mullerian duct syndrome, type II	261550
<i>ANOS1</i>	6211	Anosmin 1	Hypogonadotropic hypogonadism 1 with or without anosmia (Kallmann syndrome 1)	308700
<i>ARHGAP5</i>	675	Rho GTPase activating protein 5		
<i>ARHGAP35</i>	4591	Rho GTPase activating protein 35		
<i>AXL</i>	905	AXL receptor tyrosine kinase		
<i>CCDC141</i>	26821	Coiled-coil domain containing 141		
<i>CHD7</i>	20626	Chromodomain helicase DNA binding protein 7	Hypogonadotropic hypogonadism 5 with or without anosmia	612370
			CHARGE syndrome	214800
<i>CPE</i>	2303	Carboxypeptidase E	BDV syndrome	619326
<i>DCC</i>	2701	DCC netrin 1 receptor	Colorectal cancer, somatic	114500
			Esophageal carcinoma, somatic	133239
			Gaze palsy, familial horizontal, with progressive scoliosis, 2	617542
			Mirror movements 1 and/or agenesis of the corpus callosum	157600
<i>DLG2</i>	2901	Discs large MAGUK scaffold protein 2		
<i>DMXL2</i>	2938	Dmx like 2	Deafness, autosomal dominant 71*	617605
			Polyendocrine-polyneuropathy syndrome*	616113
			Developmental and epileptic encephalopathy 81	618663
<i>DUSP6</i>	3072	Dual specificity phosphatase 6	Hypogonadotropic hypogonadism 19 with or without anosmia	615269
<i>FEZF1</i>	22788	FEZ family zinc finger 1	Hypogonadotropic hypogonadism 22, with or without anosmia	616030
<i>FGF8</i>	3686	Fibroblast growth factor 8	Hypogonadotropic hypogonadism 6 with or without anosmia	612702

Table 1. Continued				
Gene	HGNC ID	Approved name	OMIM phenotype	Phenotype MIM number
<i>FGF17</i>	3673	Fibroblast growth factor 17	Hypogonadotropic hypogonadism 20 with or without anosmia	615270
<i>FGFR1</i>	3688	Fibroblast growth factor receptor 1	Hypogonadotropic hypogonadism 2 with or without anosmia	147950
			Encephalocraniocutaneous lipomatosis, somatic mosaic	613001
			Hartsfield syndrome	615465
			Jackson-Weiss syndrome	123150
			Osteoglophonic dysplasia	166250
			Pfeiffer syndrome	101600
			Trigonocephaly 1	190440
<i>FLRT3</i>	3762	Fibronectin leucine rich transmembrane protein 3	Hypogonadotropic hypogonadism 21 with anosmia	615271
<i>FSHB</i>	3964	Follicle stimulating hormone subunit beta	Hypogonadotropic hypogonadism 24 without anosmia	229070
<i>GNRH1</i>	4419	Gonadotropin releasing hormone 1	Hypogonadotropic hypogonadism 12 with or without anosmia*	614841
<i>GNRHR</i>	4421	Gonadotropin releasing hormone receptor	Hypogonadotropic hypogonadism 7 without anosmia	146110
<i>HESX1</i>	4877	HESX homeobox 1	Growth hormone deficiency with pituitary anomalies	182230
			Pituitary hormone deficiency, combined, 5	182230
			Septooptic dysplasia	182230
<i>HS6ST1</i>	5201	Heparan sulfate 6-O-sulfotransferase 1	Hypogonadotropic hypogonadism 15 with or without anosmia	614880
<i>IGSF10</i>	26384	Immunoglobulin superfamily member 10		
<i>IL17RD</i>	17616	Interleukin 17 receptor D	Hypogonadotropic hypogonadism 18 with or without anosmia	615267
<i>IRF2BPL</i>	14282	Interferon regulatory factor 2 binding protein like	Neurodevelopmental disorder with regression, abnormal movements, loss of speech, and seizures	618088
<i>KISS1</i>	6341	KiSS-1 metastasis suppressor	Hypogonadotropic hypogonadism 13 with or without anosmia*	614842
<i>KISS1R</i>	4510	KISS1 receptor	Hypogonadotropic hypogonadism 8 with or without anosmia	614837
			Precocious puberty, central, 1*	176400
<i>KLB</i>	15527	Klotho beta		
<i>LEP</i>	6553	Leptin	Obesity, morbid, due to leptin deficiency	614962
<i>LEPR</i>	6554	Leptin receptor	Obesity, morbid, due to leptin receptor deficiency	614963
<i>LHB</i>	6584	Luteinizing hormone subunit beta	Hypogonadotropic hypogonadism 23 with or without anosmia	228300
<i>NDNF</i>	26256	Neuron derived neurotrophic factor	Hypogonadotropic hypogonadism 25 with anosmia	618841
<i>NHLH2</i>	7818	Nescient helix-loop-helix 2	Hypogonadotropic hypogonadism 27 without anosmia*	619755
<i>NR0B1</i>	7960	Nuclear receptor subfamily 0 group B member 1	46XY sex reversal 2, dosage-sensitive	300018
			Adrenal hypoplasia, congenital	300200
<i>NSMF</i>	29843	NMDA receptor synaptonuclear signaling and neuronal migration factor	Hypogonadotropic hypogonadism 9 with or without anosmia	614838
<i>NTN1</i>	8029	Netrin 1	Mirror movements 4	618264
<i>OTUD4</i>	24949	OTU deubiquitinase 4		

Table 1. Continued				
Gene	HGNC ID	Approved name	OMIM phenotype	Phenotype MIM number
<i>PCSK1</i>	8743	Proprotein convertase subtilisin/kexin type 1	Obesity, susceptibility to, BMIQ12	612362
			Endocrinopathy due to proprotein convertase 1/3 deficiency	600955
<i>PLXNA1</i>	9099	Plexin A1	Dworschak-Punetha neurodevelopmental syndrome	619955
<i>PLXNA3</i>	9101	Plexin A3		
<i>PLXNB1</i>	9103	Plexin B1		
<i>PNPLA6</i>	16268	Patatin like domain 6, lysophospholipase	Laurence-Moon syndrome*	245800
			Boucher-Neuhauser syndrome	215470
			Oliver-McFarlane syndrome	275400
			Spastic paraplegia 39, autosomal recessive	612020
<i>POLR3A</i>	30074	RNA polymerase III subunit A	Leukodystrophy, hypomyelinating, 7, with or without oligodontia and/or hypogonadotropic hypogonadism	607694
			Wiedemann-Rautenstrauch syndrome	264090
<i>POLR3B</i>	30348	RNA polymerase III subunit B	Charcot-Marie-Tooth disease, demyelinating, type 1I	619742
			Leukodystrophy, hypomyelinating, 8, with or without oligodontia and/or hypogonadotropic hypogonadism	614381
<i>POU6F2</i>	21694	POU class 6 homeobox 2	Wilms tumor susceptibility-5	601583
<i>PROK2</i>	18455	Prokineticin 2	Hypogonadotropic hypogonadism 4 with or without anosmia	610628
<i>PROKR2</i>	15836	Prokineticin receptor 2	Hypogonadotropic hypogonadism 3 with or without anosmia	244200
<i>RAB18</i>	14244	RAB18, member RAS oncogene family	Warburg micro syndrome 3	614222
<i>RAB3GAP1</i>	17063	RAB3 GTPase activating protein catalytic subunit 1	Martsof syndrome 2	619420
			Warburg micro syndrome 1	600118
<i>RAB3GAP2</i>	17168	RAB3 GTPase activating non-catalytic protein subunit 2	Martsof syndrome 1	212720
			Warburg micro syndrome 2	614225
<i>RNF216</i>	21698	Ring finger protein 216	Cerebellar ataxia and hypogonadotropic hypogonadism	212840
<i>SEMA3A</i>	10723	Semaphorin 3A	Hypogonadotropic hypogonadism 16 with or without anosmia	614897
<i>SEMA3E</i>	10727	Semaphorin 3E		
<i>SEMA3F</i>	10728	Semaphorin 3F		
<i>SMCHD1</i>	29090	Structural maintenance of chromosomes flexible hinge domain containing 1	Bosma arhinia microphthalmia syndrome	603457
			Facioscapulohumeral muscular dystrophy 2, digenic	158901
<i>SOX2</i>	11195	SRY-box transcription factor 2	Microphthalmia, syndromic 3	206900
			Optic nerve hypoplasia and abnormalities of the central nervous system	206900
<i>SOX3</i>	11199	SRY-box transcription factor 3	Intellectual developmental disorder, X-linked, with isolated growth hormone deficiency	300123
			Panhypopituitarism, X-linked	312000
<i>SOX10</i>	11190	SRY-box transcription factor 10	PCWH syndrome	609136
			Waardenburg syndrome, type 2E, with or without neurologic involvement	611584
			Waardenburg syndrome, type 4C	613266
<i>SOX11</i>	11191	SRY-box transcription factor 11	Intellectual developmental disorder with microcephaly and with or without ocular malformations or hypogonadotropic hypogonadism	615866

Table 1. Continued				
Gene	HGNC ID	Approved name	OMIM phenotype	Phenotype MIM number
<i>SPRY4</i>	15533	Sprouty RTK signaling antagonist 4	Hypogonadotropic hypogonadism 17 with or without anosmia	615266
<i>SRA1</i>	11281	Steroid receptor RNA activator 1		
<i>STUB1</i>	11427	STIP1 homology and U-box containing protein 1	Spinocerebellar ataxia 48	618093
			Spinocerebellar ataxia, autosomal recessive 16	615768
<i>TCF12</i>	11623	Transcription factor 12	Hypogonadotropic hypogonadism 26 with or without anosmia	619718
			Craniosynostosis 3	615314
<i>TAC3</i>	11521	Tachykinin precursor 3	Hypogonadotropic hypogonadism 10 with or without anosmia	614839
<i>TACR3</i>	11528	Tachykinin receptor 3	Hypogonadotropic hypogonadism 11 with or without anosmia	614840
<i>TBC1D20</i>	16133	TBC1 domain family member 20	Warburg micro syndrome 4	615663
<i>WDR11</i>	13831	WD repeat domain 11	Hypogonadotropic hypogonadism 14 with or without anosmia	614858
			Intellectual developmental disorder, autosomal recessive 78	620237

ANOS1

The first gene identified as causative for KS is *ANOS1* (31). Formerly known as *KAL1* it is located on the short arm of the X chromosome (Xp22.3) (OMIM: 300836) (32). Ten to 20 percent of males with KS carry *ANOS1* mutations or intragenic microdeletions are present (33, 34). The extracellular glycoprotein it encodes, anosmin-1, plays a role in the adhesion of GnRH cells and axon migration during organogenesis (35). Anomin-1 exerts its biological effects mainly through signal modulation of *FGFR1* via its third fibronectin-like type 3 (FnIII) domain and the N-terminal region (36). The migratory defect of olfactory and GnRH neurons is the central mechanism underlying the clinical features of *ANOS1* mutations (37). In its first observation, in a 19-week-old male human fetus with a deletion in *ANOS1*, GnRH neurons could not migrate to their normal positions in the brain (38). In KS cases associated with *ANOS1* mutations, penetrance has been reported to be almost complete (39,40). Additional clinical findings include bimanual synkinesis, unilateral renal agenesis, vas deferens agenesis, and deafness (28,41).

FGFR1, FGF8 and related genes (FGF17, IL17RD, DUSP6, SPRY4, FLRT3, and KLB)

FGFR1 encodes a receptor belonging to the tyrosine kinase superfamily. It regulates central developmental processes, such as neuronal proliferation, differentiation, and migration critical for embryonic development. *FGFR1* is the first gene in which mutations were identified for the AD form of KS (42). However, over time, *FGFR1* has also been found to be associated with nIHH (43,44). Around 10% of patients with KS were found to have inactivating mutations in *FGFR1* (29,43,45). Loss-of-function

mutations in *FGFR1* were detected in 7% of 134 nIHH patients (46). To date numerous insertions/deletions, missense, and non-sense mutations have been reported with AD, AR, *de novo*, and oligogenic inheritance (29,47,48). Loss of *FGFR1* function elicit reproductive abnormalities ranging from severe AD KS through fully penetrant nIHH to delayed puberty (43,44,45,49,50). *FGFR1* mutations have been associated with cleft palate, synkinesis, and tooth agenesis, and asymptomatic carriers have been reported in some familial cases (5,51).

FGF8 and *FGF17* are *FGFR1* ligands with similar sequence structures that play a role in GnRH neuron ontogenesis. Mutations in these genes have been reported in IHH patients with varying olfactory functions (52,53). Mice homozygous for the hypomorphic *Fgf8* allele exhibited absent olfactory bulbs and lacked GnRH neurons in the hypothalamus (52). IHH patients harboring *FGF8* variants have also been reported to exhibit additional phenotypic features, including cleft lip and/or palate, a flat nasal bridge, and camptodactyly (54,55). Further screening for *FGF8*-related genes in a group of 388 congenital IHH patients revealed inactivating variants in *FGF17*, *IL17RD*, *DUSP6*, *SPRY4*, and *FLRT3* (53).

KLB encodes β -Klotho, a co-receptor essential for *FGF21* signaling via *FGFR1*. In one study, over 300 IHH patients were screened, identifying 13 individuals with loss-of-function *KLB* variants. Most of these patients exhibited metabolic abnormalities, including insulin resistance or dyslipidemia. Notably, *Klb* knockout mice displayed a milder hypogonadal phenotype compared to the human presentation (56).

HS6ST1

heparan sulfate 6-O-sulfotransferase 1 (HS6ST1) is directly involved in the sulfation of heparan sulfate proteoglycans, which are critical modulators of FGF signaling. The 6-O-sulfation of heparan sulfate chains, catalyzed by *HS6ST1*, is required for optimal binding and activation of FGF ligands (such as *FGF8* and *FGF17*) to their receptor *FGFR1* (57). This interaction is essential for the development, migration, and survival of GnRH neurons during embryogenesis. Mutations in *HS6ST1*, often co-occurring with variants in other known KS genes, have been reported in seven families (58).

PROKR2 and PROK2

PROK2 and *PROKR2* encode prokineticin 2, an 81-amino acid peptide, and its G protein-coupled receptor, respectively. Both play critical roles in the development of neuronal precursors and are essential for processes such as olfactory bulb morphogenesis and sexual maturation (59). This ligand-receptor pair has been identified as a strong candidate for the pathogenesis of KS as *PROK2* (60,61) or *PROKR2* knockout mice had defective olfactory bulbs and failed migration of GnRH neurons (62). Subsequent studies identified inactivating variants in *PROKR2* and *PROK2* in patients with KS. The majority of these mutations have been identified in the heterozygous state, although both homozygous and compound heterozygous variants have also been reported (63). Patients with *PROK2* or *PROKR2* mutations have considerable phenotypic variability (61,64,65), ranging from KS to nIHH. A variety of associated clinical features has been reported in affected individuals, including fibrous dysplasia, synkinesia, epilepsy, and Crohn's disease (66). Mutations in *PROKR2* and *PROK2* are frequently identified in combination with variants in other genes, supporting an oligogenic mode of inheritance in IHH.

CHD7

CHARGE syndrome is a multisystem disorder that includes Coloboma, Heart anomalies, choanal Atresia, growth Retardation, Genital defects and Ear anomalies (67). *CHD7* mutations are present in the majority of patients with CHARGE syndrome. *CHD7* is a chromatin-remodeling protein essential for the ontogeny of GnRH neurons and the proper targeting of olfactory axons during embryogenesis. Pathogenic *CHD7* mutations disrupt these developmental processes, resulting in a reduced number of hypothalamic GnRH neurons and defective GnRH secretion. There is a range of abnormalities in the GnRH neuron migration pathway in mice with *CHD7* deficiency (68,69). While large *de novo* deletions are noted in classical CHARGE syndrome patients, inherited or *de novo* point mutations may result in KS/nIHH (48,70,71). Thus, IHH patients should be carefully examined for possible clinical features of CHARGE syndrome, such as abnormal ears, deafness, semicircular canal hypoplasia, and coloboma (67).

WDR11

WDR11 in partnership with EMX1, a homeodomain transcription factor, is essential for normal Hedgehog (Hh) signaling and ciliogenesis, both of which are critical for the embryonic development and migration of GnRH neurons. Mutations in *WDR11* disrupt Hh pathway signaling, impairing the formation and function of primary cilia, which are required for the proper migration of GnRH and olfactory neurons (72). By positional cloning, heterozygous mutations were discovered in several patients with KS (73).

CCDC141

CCDC141 encodes a coiled-coil domain containing protein that is expressed in GnRH neurons. We have reported inactivating *CCDC141* variants in four separate families with IHH. Affected individuals exhibit normal olfaction and anatomically normal olfactory bulbs (74). In a rodent nasal explant model, knockdown of *Ccdc141* led to impaired embryonic migration of GnRH neurons without affecting olfactory axon outgrowth, thereby producing a nIHH phenotype distinct from other genes implicated in GnRH neuronal migration (7). *CCDC141* Mutations have been identified as a recurrent finding in individuals with CDGP. Among a cohort of 193 patients with CDGP, 21 individuals (6%) were found to carry predicted deleterious variants in *CCDC141* (75).

FEZF1

The protein product of *FEZF1* facilitates the penetration of olfactory receptor neuron axons through the basal lamina of the central nervous system in murine models. As a subset of these axons serves as a migratory scaffold for GnRH neurons, *FEZF1* deficiency results in failed entry of GnRH neurons into the brain (76,77). Through autozygosity mapping and WES of 30 individuals with KS, we discovered homozygous loss-of-function mutations in *FEZF1* in two separate consanguineous families, each with two affected siblings (78). *FEZF1* mutations are apparently extremely rare as no new KS cases have been reported to date.

IGSF10

IGSF10, a member of the immunoglobulin superfamily, was implicated in delayed puberty by Howard et al. (8), who analyzed WES data from over 100 affected individuals and identified pathogenic mutations in six families. Knockdown studies demonstrated reduced GnRH neuronal migration in the GN11 cell line. Despite this impaired migration, patients harboring *IGSF10* mutations exhibited a normal sense of smell. The authors proposed that a reduced number or delayed arrival of GnRH neurons to the hypothalamus results in a milder disruption of the GnRH neuronal network, manifesting as delayed puberty rather than permanent IHH. Notably, *IGSF10* mutations were also identified in adults with functional hypothalamic amenorrhea, a condition considered a mild and reversible form of HH (8).

SEMA3A and related genes (*SEMA3E*, *SEMA3G*, *SEMA3F*, *PLXNA1*, *PLXNA3* etc)

The precise targeting of GnRH neurons and olfactory/vomer nasal projections relies on the coordinated activity of axonal guidance cues, including semaphorins which are a large and heterogeneous family of secreted and membrane-bound proteins (79). Mutations in class-3 semaphorin family members, including *SEMA3A*, *SEMA3E*, and *SEMA3G*, have been implicated in the pathogenesis of IHH (80,81,82). *SEMA3* proteins exert their biological functions by binding to Neuropilin co-receptors, forming heteromeric complexes with *PlexinA1-4* (*PLXNA1-4*) receptors, thereby initiating plexin-mediated signal transduction pathways (83). Non-synonymous heterozygous variants in *PLXNA1* have been identified in KS individuals (84). More recently we have identified deleterious variants in *SEMA3F* and *PLXNA3* that caused IHH (85).

SEMA3F and PLXNA3

SEMA3F and its coreceptor *PLXNA3* play a role in cell migration and axonal guidance (86). WES of 216 patients with IHH identified rare *SEMA3F* and *PLXNA3* variants in 15 individuals. Over half (54%) also carried mutations in other known IHH genes, highlighting the disorder's oligogenic nature. *SEMA3F* variants followed AD inheritance with variable penetrance, while *PLXNA3* variants were X-linked recessive. Six patients exhibited impaired olfaction. The study provided clinical, genetic, and cellular evidence supporting the role of *SEMA3F* signaling deficiency in IHH pathogenesis (85).

PLXNA1

Plexin-A1, a transmembrane coreceptor for semaphorin 3 signaling, is encoded by *PLXNA1* (87). Heterozygous *PLXNA1* variants were identified in 15 of 237 unrelated patients with KS, and impaired plexin-A1 signaling has been linked to oligogenic inheritance in KS (84). Subsequently, by screening the WES data of 215 IHH patients, we identified rare heterozygous *PLXNA1* variants in KS and nIHH patients carrying additional variants in known IHH genes. Thus, the contribution of *PLXNA1* to the oligogenicity of both forms of IHH was confirmed (88).

PLXNB1

The receptor for semaphorin 4D, plays a critical role in GnRH neuronal development. In murine models, disruption of *Sema4D/PLXNB1* signaling results in abnormal GnRH ontogeny. In a cohort of 336 patients with IHH, we were able to detect six rare *PLXNB1* variants in eight individuals with the nIHH (89).

SMCHD1

SMCHD1 encodes an epigenetic repressor that is expressed in the human olfactory epithelium. Shaw et al. (90) identified inactivating mutations in *SMCHD1* as the underlying cause

of congenital arhinia in 41 cases. Notably, 97% of affected individuals also exhibited hypogonadal features, including cryptorchidism, microphallus, or amenorrhea, alongside absent olfactory structures and anosmia (90).

SOX10

Inactivating mutations in *SOX10* are responsible for Waardenburg syndrome, a rare condition marked by pigmentation defects and sensorineural hearing loss. In a subset of KS patients presenting with deafness, *SOX10* mutations were identified in about one-third of cases. Consistent with these findings, *Sox10* knockout mice exhibit a complete absence of olfactory ensheathing cells along the olfactory nerve pathway, highlighting the critical role of this gene in olfactory system development (91). A large cohort study of 1309 IHH patients reported that developmental problems due to *SOX10* variants may encompass a phenotypic line from KS to nIHH (92).

SOX2

SOX2 encodes the SRY-related, HMG-box 2 transcription factor protein. A study involving eight IHH patients with heterozygous *SOX2* variants who had severe eye defects found that pathogenic *SOX2* variants were linked to both anosmic and normosmic forms of IHH. Functional analyses indicated that *Sox2* was highly expressed in the hypothalamus of adult mice. The study emphasized that screening for *SOX2* variants should be performed in patients, regardless of the presence of ocular defects, when conducting genetic evaluations for IHH (93).

NDNF

NDNF is a secreted neurotrophic factor involved in the migration of GnRH neurons and is a member of the fibronectin type III (FN3) superfamily. Screening for rare variants in FN3 domain-containing proteins identified three protein-truncating and one missense heterozygous *NDNF* variant among patients with KS. In *NDNF*-null mice, a reduced number of GnRH neurons reached their final destination compared to wild-type *NDNF^{F+/+}* mice (94). More recently, we identified a homozygous protein-truncating variant in *NDNF* in a consanguineous family with KS, highlighting that, in addition to the previously described dominant inheritance, *NDNF*-related disease can also follow an AR pattern (95).

AMH and AMHR2

Anti-Müllerian Hormone (AMH) is expressed in migrating GnRH neurons in mouse and human fetuses during embryonic development and functions as a promotility factor (96). *AMHR2*-deficient mice exhibit aberrant development of the peripheral olfactory system and impaired embryonic migration of GnRH neurons. In humans, heterozygous inactivating variants in *AMH* or *AMHR2* have been associated with IHH. These findings indicate

the critical role of *AMH/AMHR2* signaling in GnRH neuronal migration and its contribution to the pathogenesis of IHH (97).

AXL

AXL receptor tyrosine kinases, members of the TAM (TYRO3/*AXL/MERTK*) family, play a role in GnRH neuron migration and survival. Studies of sexual maturation in *AXL* null mice reported that TAM function was impaired (98). *AXL* variants have been identified in both anosmic and normosmic IHH probands.

NTN1

Netrin-1, encoded by the *NTN1* gene, plays a crucial role in central nervous system development by guiding axonal and neuronal migration through its receptor DCC. In *DCC*^{-/-} and *NTN1*^{-/-} mouse embryos, GnRH neurons exhibited aberrant trajectories and failed to reach the medial preoptic area, highlighting the importance of NTN1/DCC signaling in proper GnRH neuronal migration (99,100). WES of a cohort of 133 individuals with IHH identified pathogenic variants in *NTN1* and the gene for its receptor, *DCC*. Five heterozygous *DCC* variants were detected in six probands, five of whom had KS and one with nIHH. In addition, co-occurring variants in both *DCC* and *NTN1* were identified in two KS patients, supporting an oligogenic basis for disease pathogenesis (101).

nIHH Associated Genes

Genes implicated in nIHH are particularly informative for understanding the regulation of the HPG axis and the timing of puberty. Genetic analyses of familial nIHH cases have significantly advanced this understanding. In a study of 22 consecutive multiplex families with nIHH, mutations were identified in five genes, *GNRHR*, *TACR3*, *TAC3*, *KISS1R*, and *KISS1*, in 77% of families. Among these, *GNRHR* and *TACR3* mutations were the most frequently observed, each accounting for approximately one-third of the genetically resolved cases (24).

GNRHR and GNRH1

GNRH1 and *GNRHR* are the most obvious candidate gene in the etiology of IHH. In 1997, de Roux et al. (102) identified compound heterozygous mutations in *GNRHR* in two siblings with partial nIHH, showing that Gln106Arg impaired GnRH binding while Arg262Gln reduced IP3 signaling. The male sibling exhibited normal gonadotropin levels and LH pulse frequency but reduced pulse amplitude, consistent with partial GnRH receptor dysfunction. Shortly after, Layman et al. (103) reported a family with four siblings carrying compound heterozygous *GNRHR* mutations (p.Arg262Gln and p.Tyr284Cys), further supporting the role of biallelic *GNRHR* mutations in IHH without anosmia or developmental anomalies (103). Subsequent studies found *GNRHR* variants in approximately 5-6% of nIHH cases (104). This relatively high prevalence of *GNRHR* was confirmed

in subsequent studies (105). To date over 60 distinct mutations have since been reported (106).

Genotype-phenotype correlations have been observed for specific *GNRHR* mutations. The genetic makeup (homozygous, compound heterozygous, or monoallelic variants) broadly correlates with clinical severity, ranging from complete IHH to milder forms, such as CDGP and functional hypothalamic amenorrhea (104). The homozygous R139C missense mutation in the conserved DRS motif of the GnRH receptor causes complete IHH by severely impairing receptor trafficking to the plasma membrane, a defect reversible with the pharmacological chaperone IN3 (107). In contrast, the heterozygous Gln106Arg mutation is linked to adult-onset IHH (AOHH), where normal pubertal development precedes nIHH. Homozygosity for p.Gln106Arg has also been associated with the fertile eunuch variant of nIHH, characterized by hypogonadism with preserved testicular size and partial virilization (108). These findings highlight how specific *GNRHR* mutations contribute to a broad spectrum of GnRH deficiency phenotypes (5,106).

GNRH1 encodes the GnRH preprohormone. Deletion of *GNRH1* in murine models was shown to result in complete absence of GnRH synthesis, a finding reported well before analogous mutations were identified in humans (109,110). Over a decade after the initial discovery of *GNRHR* mutations, pathogenic *GNRH1* variants were reported in humans (111,112). Bouligand et al. (111) demonstrated that pulsatile GnRH administration for two weeks resulted in synchronous LH pulses, increased levels of estradiol, and a single dominant ovarian follicle. These findings confirmed the hypothalamic origin and pivotal role of GnRH in human reproduction. Affected individuals frequently present with micropenis and cryptorchidism (111,112,113).

KISS1R and KISS1

In 2003, kisspeptin emerged as a pivotal central regulator of GnRH neuronal activity following the identification of mutations in a previously little-characterized G protein-coupled receptor, initially termed *GPR54* and later renamed *KISS1R* (kisspeptin receptor) (106). In 2003, two independent research groups concurrently reported homozygosity mapping in familial cases of IHH, resulting in the first identification of pathogenic mutations in *KISS1R* (114,115). Mutant *KISS1R* constructs exhibited impaired receptor function in *in vitro* assays, and *KISS1R*-knockout mice recapitulated the human hypogonadotropic phenotype, confirming the essential role of the kisspeptin signaling pathway in pubertal and reproductive regulation across mammals (114). In a mutational screening study, only five out of 166 (3%) probands with nIHH were found to have rare variants in *KISS1R* (116). The rarity of mutations in *KISS1* and *KISS1R* may be attributed to evolutionary selection pressures, given the critical roles of kisspeptin in placentation, reproductive

function, and metastasis suppression, which likely constrain the transmission of deleterious variants within populations (106). Studying a large, consanguineous family with four sisters with nIHH, we found inactivating mutations altering the 4th amino acid of Kisspeptin-10. Overnight frequent LH sampling did not reveal any LH pulsatility, further confirming the essential role of kisspeptin signaling in the GnRH pulse generator (117). Coutant et al. (118) recently identified homozygous frameshift mutations in *KISS1* within a consanguineous family (118). Molecular analyses confirmed a complete absence of kisspeptin protein. Affected male siblings exhibited congenital gonadotropin deficiency, including bilateral cryptorchidism, micropenis, and absent spontaneous puberty. However, the two older brothers later showed spontaneous reversal of hypogonadism, with normalization of testicular volume and spermatogenesis. These findings indicate that complete kisspeptin deficiency does not preclude delayed GnRH activation or pubertal maturation, highlighting the redundancy and adaptability of upstream neuroendocrine pathways (118). The potential involvement of alternative *KISS1R* ligands, neuroendocrine plasticity, or compensatory pathways, such as neurokinin B (NKB) or glutamatergic signaling, requires further investigation.

TACR3 and TAC3

Tachykinin receptor 3, encoded by *TACR3*, mediates the biological actions of neurokinin B (NKB), which is encoded by *TAC3*. Through autozygosity mapping aimed at discovering novel regulators of the HPG axis, we identified homozygous non-synonymous mutations in the coding regions of *TAC3* or *TACR3* in nine individuals from four families presenting with an nIHH phenotype (119). With the additional cases identified in our cohort, it became clear that *TACR3* mutations are almost as common as *GNRHR* mutations (24,120). Similar findings regarding the prevalence of *TACR3* mutations have been reported by other research groups. Gianetti et al. (121) found 19 among 345 (5.5%) cases while a very similar rate (5.2%) was observed by Francou et al. (122) from a cohort of 173 cases of familial and sporadic nIHH. The frequent occurrence of micropenis and cryptorchidism in male patients with *TACR3* mutations suggests that functional *TACR3* signaling is essential for normal fetal gonadotropin secretion, which in turn regulates testicular development, descent, and penile growth (4).

Clinical reversibility, characterized by spontaneous pubertal progression, often following a period of exogenous sex steroid therapy, was observed in approximately 10% of an unselected cohort with nIHH (12). Gianetti et al. (121) reported a significantly higher rate of reversibility of 83% in their cohort of patients with *TAC3/TACR3* mutations. In our cohort, four patients from three unrelated and ethnically diverse families exhibited clinical recovery, representing 25% (4/16) of the cases. Notably, all of

these families carried the same *TACR3* mutation (p.Thr177Lys). Given the relatively high rate of reversibility, it was hypothesized that CDGP might represent a mild form of IHH linked to *TACR3* mutations. To investigate this, Vaaralahti et al. (123) screened *TAC3* and *TACR3* in 146 Finnish individuals with CDGP but identified no pathogenic variants associated with the phenotype.

Additional clinical studies have enhanced our understanding about the regulation of the HPG axis by Neurokinin B signaling. Young et al. (124) showed that patients with null mutations in *TAC3* could achieve pubertal levels of gonadotropins and sex steroids following repeated administration of exogenous GnRH. This finding indicates that neurokinin B acts at a hypothalamic level, upstream of GnRH secretion, rather than directly influencing pituitary function (124). Furthermore, a genome-wide association study identified a significant association between age at menarche, a surrogate marker of pubertal start, and a single nucleotide polymorphism (rs3733631) located immediately upstream of *TACR3*, supporting a role for neurokinin B signaling in the regulation of pubertal timing at the population level (125).

IRF2BPL (EAP1)

Pubertal onset is postulated to be regulated in part by transcriptional factors such as *EAP1* (126). In a cohort with familial CDGP, two rare *EAP1* variants (p.Ala221del and p.Asn770His) were identified, both impairing GnRH promoter activation through distinct molecular mechanisms. These findings provide the first link between *EAP1* mutations and CDGP (127).

LEP and LEPR

Leptin deficiency with mutations in either *LEP* (encoding leptin) or *LEPR* (encoding the leptin receptor) is associated with IHH (128,129). Administration of leptin in individuals with *LEP* deficiency restores normal pubertal development but does not induce precocious puberty in prepubertal children. This suggests that leptin functions as a permissive, rather than initiatory, factor in the onset of puberty in humans (130). These patients are readily distinguishable from other individuals with IHH due to the characteristic presentation of early-onset obesity and hyperphagia.

NROB1 (DAX1)

NROB1 belongs to the nuclear receptor superfamily and is classified as an orphan receptor due to the absence of a known endogenous ligand. Mutations in *NROB1* are known to cause adrenal hypoplasia congenita in combination with IHH (131,132,133,134,135). Adrenal hypoplasia typically presents as adrenal insufficiency during infancy, whereas IHH becomes manifest in affected males who survive into the second decade of life. Nuclear receptors, such as SF-1 and LRH-1, involved in

adrenal and gonadal physiology and development, are regulated in their transcriptional activity by coregulatory molecules (136). *DAX-1*, lacks a DNA-binding domain and functions exclusively as a coregulator (137). Notably, *Dax-1* is predominantly expressed in the arcuate nucleus (ARC) of the hypothalamus. In adult female mice, *DAX-1* is present in at least 70% of *Kiss1* neurons within the ARC, which is associated with pubertal development, whereas it is found in fewer than 5% of *Kiss1* neurons in the AVPV nucleus, which is exclusively linked to the menstrual cycle (138). These findings suggest that *DAX-1* is selectively involved in the regulation of pubertal onset and the sustained function of the HPG axis. As mentioned earlier, mutations in *NROB1* result in adrenal hypoplasia congenita together with IHH (131,132). Paradoxically, *NROB1* mutations can also result in the opposite phenotype, precocious puberty, even within the same kindred (133,134,135). The genetic mechanisms underlying these divergent phenotypic outcomes remain poorly understood, highlighting intriguing genotype-phenotype correlations. This paradox suggests a complex, context-dependent role for *DAX-1* in regulating the HPG axis and pubertal timing.

SRA1

SRA1 was the first gene demonstrated to exert its function through both its protein product and a non-coding, functional RNA transcript (139). These proteins serve as co-regulators for nuclear receptors, including sex steroid receptors, and play a critical role in modulating the activity of SF-1 and LRH-1, the principal regulators of steroid hormone biosynthesis. *SRA1* is required for the synergistic enhancement of SF-1 transcriptional activity by *DAX-1* (*NROB1*), mutations in which also cause IHH (140). We and others have reported nIHH patients with inactivating *SRA1* mutations (141,142,143).

PNPLA6

Gordon Holmes syndrome (GDHS) is characterized by cerebellar ataxia/atrophy and nIHH, while the related Boucher-Neuhäuser syndrome also includes chorioretinal dystrophy. *PNPLA6*, which encodes neuropathy target esterase (NTE), a key regulator of phospholipid metabolism, was found to carry loss-of-function mutations in six GDHS patients from three unrelated families via autozygosity mapping and WES. Functional studies showed that NTE inhibition in β 2 gonadotroph cells impaired LH exocytosis in response to GnRH. These findings suggest that NTE dysfunction in GDHS disrupts phospholipid homeostasis, contributing to both neurodegeneration and impaired LH secretion, resulting in nIHH (144).

OTUD4 and RNF216

Ubiquitination-related *OTUD4* encodes a deubiquitinase, while *RNF216* encodes a ubiquitin E3 ligase. *OTUD4* and *RNF216* mutations have been identified in patients with GDHS.

Patients have progressive ataxia, dementia, and neuronal losses are observed in the cerebellar pathway and hippocampus. Functional studies have shown that knockout of *OTUD4* and *RNF216* in zebrafish causes defects in the eye and cerebellum and that suppression of the two genes together worsens these phenotypes. Hence, inactivating mutations in *OTUD4* and *RNF216* cause neurodegeneration and reproductive failure through dysregulated ubiquitination (145).

STUB1

STUB1 encodes C-terminus of HSC70-inactivating protein, which functions as a E3 ubiquitin ligase. Pathogenic variants of *STUB1* have been associated with GDHS (146).

POU6F2

POU6F2 belongs to a gene family characterized by a bipartite DNA-binding domain, comprising a POU-specific domain and a POU homeodomain. Members of this family function as transcriptional regulators involved in cell type-specific differentiation. Several POU domain-containing proteins have been implicated in the regulation of GnRH neuron expression (147,148). Using WES data from two independent IHH cohorts (331 nIHH, 85 KS; 416 patients in total and 677 nIHH, 632 KS; 1309 patients in total), 12 rare missense variants of *POU6F2* were identified in 15 patients. Functional studies of two different isoforms encoded by *POU6F2* were performed, and the function of isoform 1 was proven as a transcriptional regulator of *GNRH1* expression. Thus, pathogenic *POU6F2* variants were shown to be involved in IHH pathogenesis by disrupting normal GnRH migration (149).

DLG2

DLG2 encodes a scaffolding protein that interacts with N-methyl-D-aspartate (NMDA) receptors, which have been implicated in the regulation of sexual maturation in animal models. WES identified a rare missense variant in *DLG2* in a large family with delayed puberty. Functional studies demonstrated that this variant reduces *GnRH* expression *in vitro*, suggesting a potential mechanistic link between *DLG2* and pubertal timing (150). A subsequent study screened the WES data of 336 IHH probands from 290 independent families for rare *DLG2* variants. A total of one homozygous and two heterozygous missense variants were identified in three independent normosmic patients (151).

NHLH2

NHLH2, a basic helix-loop-helix transcription factor family member, mediates leptin-induced activation of POMC in the leptin-melanocortin pathway. Screening of WES data in a large IHH cohort revealed obese patients with rare disease-causing sequence variants. *In silico* and *in vitro* analyses of the findings showed that *NHLH2* binding to the *Mc4r* promoter

and *KISS1* transactivation were reduced, supporting a critical role for *NHLH2* in human puberty and body weight control (152). Remarkably, *NHLH2* knockout mice exhibit a phenotype closely resembling that of patients with rare inactivating *NHLH2* variants, characterized by nIHH and late-onset obesity (153).

CPE

CPE encodes an enzyme responsible for processing neuropeptides, including GnRH, into their biologically active forms within the hypothalamus. Inactivating mutations in *CPE* result in a syndrome characterized by severe obesity, intellectual disability, disrupted glucose homeostasis, and IHH, a phenotype consistent with observations in *CPE* knockout mouse models (154). A subsequent study detected a homozygous non-sense *CPE* mutation in three obese siblings with mental retardation and IHH (155). Comparison with previously reported cases led to the delineation of a distinct clinical entity termed Blakemore-Durmaz-Vasileiou (BDV) syndrome, which is an extremely rare AR disorder characterized by a combination of impaired intellectual development, hyperphagia, and IHH (156).

POLR3A and POLR3B

RNA polymerase III regulates fundamental cellular processes through the transcription of small RNAs. Its catalytic core is composed of multiple subunits, including *POLR3A* and *POLR3B*. Pathogenic variants in these subunits have been associated with 4H syndrome (also known as POLR3-related leukodystrophy), a rare disorder characterized by hypomyelination, hypodontia, and IHH (157,158). Mice studies have shown that missense mutations in *POLR3A* and *POLR3B* can variably disrupt development and Pol III function (159). It is still unclear how inactivating mutations in those genes cause IHH.

Small GTPase related genes (*RAB18*, *RAB3GAP1*, *RAB3GAP2*, *TBC1D20*, and *DMXL2*)

Mutations in several genes related to small GTPases that include *RAB18*, *RAB3GAP1*, *RAB3GAP2*, *TBC1D20*, and *DMXL2*, have been implicated in IHH, often in conjunction with neurodegenerative features. Small GTPases are critical regulators of intracellular trafficking, particularly in endocytosis and exocytosis. *RAB18* is a member of the Ras-related GTPases that play a role in apical endocytosis/recycling between the plasma membrane and early endosomes (160). Mutations in *RAB18* or in any of its essential regulators, *RAB3GAP1*, *RAB3GAP2*, and *TBC1D20* (161,162,163), are associated with Warburg micro syndrome type 3 (164). Warburg micro and Martsolf syndromes are overlapping clinical entities characterized by IHH, progressive spasticity, severe developmental delay, microcephaly, cortical visual impairment, hypotonia, and optic nerve atrophy. *DMXL2* encodes for rabconnectin-3a, which is a regulator of another intracellular GTPase, *RAB3A*. Rabconnectin-3a is expressed in exocytosis vesicles in GnRH axons in the median eminence of the hypothalamus (165). Furthermore, inactivating

DMXL2 mutations cause a novel complex syndrome that features IHH and a neurodegenerative phenotype, including cerebellar ataxia and demyelinating polyneuropathy, among other clinical features (165).

ARHGAP35 and ARHGAP5

ARHGAP35 which codes for Rho GTPase activating protein 35 and *ARHGAP5* which codes for Rho GTPase activating protein 5 are Rho GTPase activating protein genes. Rare protein-truncating variants (PTVs) in *ARHGAP35* have been reported to result in IHH. Zebrafish modeling has shown that neuronal areas are reduced in mutant embryos lacking the *ARHGAP35* paralog *ARHGAP35*. No changes were observed in the *ARHGAP35* paralog in functional studies, and it was identified as an IHH candidate. These observations suggest a novel role for the p190 RhoGAP proteins in GnRH neuronal development and integrity (166).

FSHB

FSHB encodes the beta subunit of follicle-stimulating hormone (FSH). A homozygous deletion of *FSHB* has been reported in a patient with nIHH, primary amenorrhea, and infertility due to isolated pituitary FSH deficiency (167). Studies have reported mutations in compound heterozygous, missense, and non-sense types (168,169). Mouse studies show that *FSHB*^{-/-} female mice are sterile and hypogonadal (170).

LHB

Luteinizing hormone (LH), encoded by *LHB*, is a glycoprotein hormone essential for the regulation of gonadal function. A homozygous mutation in *LHB* was first identified in a patient with nIHH caused by biologically inactive LH (171,172). Subsequent studies have reported missense mutations, non-sense mutations, and small deletions in *LHB* associated with nIHH (173,174,175). In animal models, targeted disruption of *LHB* in mice resulted in reduced testicular size and decreased testosterone levels in males, while females exhibited a hypogonadal phenotype (176).

Scientific Significance of Identifying Ihh-Associated Genes

Undoubtedly, the most impactful contribution of IHH-associated gene studies has been the elucidation of the long-sought GnRH pulse generator, advancing our fundamental understanding of reproductive neuroendocrine regulation (177,178,179,180). A surge of research into kisspeptin and neurokinin B signaling, catalyzed by the discovery of inactivating mutations in familial cases of nIHH, has led to the characterization of the elusive GnRH pulse generator. Current understanding centers on a population of sex steroid-responsive neurons within the arcuate (infundibular) nucleus that co-express Kisspeptin, NKB, Dynorphin (KNDy), and estrogen receptor alpha (ER α), collectively termed KNDy neurons. Within this network, stimulatory signals from NKB initiate action potentials, which are subsequently attenuated by inhibitory dynorphin signaling. When dynorphin-mediated inhibition is overcome, a new cycle of NKB-induced excitation

ensues, resulting in rhythmic, intermittent action potentials. Each burst drives pulsatile kisspeptin release onto GnRH neuron terminals in the median eminence, thereby triggering GnRH secretion into the portal circulation and ultimately stimulating pituitary gonadotropes. The synchronization of KNDy neuronal activity is thought to be mediated by NKB-neurokinin-3 receptor (NK3R) signaling via ipsilateral and contralateral projections within the KNDy network (178,181,182).

Clinical Significance of Identifying IHH-Associated Genes

A major impact stemming from IHH-associated gene studies may be the translation into new therapeutic modalities. The first therapeutic opportunities linked to the identification of IHH genes stemmed from the discovery of *TAC3* and *TACR3* mutations in patients with nIHH (183). Antagonism of neurokinin B signaling has been used in the development of pharmacological therapies targeting two of the most common reproductive health disorders in women globally: menopausal hot flashes and polycystic ovary syndrome (PCOS).

In menopausal women, the decline in ovarian estrogen levels reduces negative feedback on KNDy neurons, causing them to become hypertrophied and to overproduce neurokinin B. KNDy neurons project to the *TACR3*-expressing median preoptic nucleus within the hypothalamus, a key region involved in thermosensory processing and heat-defense mechanisms (184,185). Building on these observations, the Rance laboratory demonstrated that ablation of KNDy neurons in rats leads to a reduction in tail-skin temperature, indicating that NKB promote cutaneous vasodilation, a key physiological component of hot flashes (186). Following clinical trials, fezolinetant, which is a selective *NK3R* antagonist, was approved for the treatment of vasomotor symptoms in menopausal women in 2023 (187,188).

NK3R antagonists also have potential for the treatment of PCOS. In premenopausal women, *NK3R* antagonism decreases the GnRH pulse frequency leading to reduced basal LH secretion, lower LH/FSH ratio, and the modulation of the temporal dynamics of ovarian sex hormone production over the menstrual cycle (189). The *NK3R* antagonist, MLE4901, was demonstrated to reduce LH pulse frequency, as well as serum LH and testosterone levels, in women with PCOS (190). These hormonal findings were validated in a recent study involving fezolinetant. However, no significant improvement was observed in menstrual cycle regularity or clinical outcome scores (187). The investigators noted that the 12-week treatment duration in this trial may have been insufficient to elicit measurable changes, as favorable clinical outcomes in PCOS trials are typically observed after 6 to 9 months of therapy (187,191). The use of an *NK3R* antagonist as a therapeutic agent for PCOS remains a promising strategy, given its potential to modulate the neuroendocrine dysregulation underlying the condition.

Concluding Remarks

Currently, approximately half of the genes underlying IHH remain unidentified. The complexity of genotype-phenotype correlations in IHH, largely due to the established phenomena of oligogenic inheritance and spontaneous or treatment-induced clinical reversibility, poses significant challenges to gene discovery. Nevertheless, advances in next-generation sequencing technologies are expected to drive continued progress in uncovering the genetic basis of IHH. These investigations not only enhance our understanding of fundamental biological processes, such as the recent elucidation of the GnRH pulse generator, but also inform the development of targeted therapeutics, exemplified by the approval of fezolinetant for the treatment of menopausal hot flashes.

Footnotes

Authorship Contributions

Surgical and Medical Practices: A. Kemal Topaloğlu, Concept: A. Kemal Topaloğlu, Design: A. Kemal Topaloğlu, Data Collection or Processing: A. Kemal Topaloğlu, Analysis or Interpretation: A. Kemal Topaloğlu, Literature Search: A. Kemal Topaloğlu, Leman Damla Kotan, Writing: A. Kemal Topaloğlu, Leman Damla Kotan.

Conflict of Interest: One author of this article, A. Kemal Topaloğlu, is a member of the Editorial Board of the Journal of Clinical Research in Pediatric Endocrinology. However, he did not involved in any stage of the editorial decision of the manuscript. The editors who evaluated this manuscript are from different institutions. The other authors declared no conflict of interest.

Financial Disclosure: The authors declared that this study received no financial support.

References

1. Duittoz AH, Forni PE, Giacobini P, Golan M, Mollard P, Negrón AL, Radovick S, Wray S. Development of the gonadotropin-releasing hormone system. *J Neuroendocrinol*. 2021:e13087. Epub 2022 Jan 23
2. Plant TM. Neuroendocrine control of the onset of puberty. *Front Neuroendocrinol*. 2015;38:73-88. Epub 2015 Apr 22
3. Saengkaew T, Howard SR. Genetics of pubertal delay. *Clin Endocrinol (Oxf)*. 2022;97:473-482. Epub 2021 Oct 13
4. Grumbach MM. A window of opportunity: the diagnosis of gonadotropin deficiency in the male infant. *J Clin Endocrinol Metab*. 2005;90:3122-3127. Epub 2005 Feb 22
5. Topaloğlu AK. Update on the genetics of idiopathic hypogonadotropic hypogonadism. *J Clin Res Pediatr Endocrinol*. 2017;9(Suppl 2):113-122. Epub 2017 Dec 27
6. Semple RK, Topaloglu AK. The recent genetics of hypogonadotropic hypogonadism - novel insights and new questions. *Clin Endocrinol (Oxf)*. 2010;72:427-435. Epub 2009 Aug 29
7. Hutchins BI, Kotan LD, Taylor-Burds C, Ozkan Y, Cheng PJ, Gurbuz F, Tiong JD, Mengen E, Yuksel B, Topaloglu AK, Wray S. *CCDC141* mutation identified in anosmic hypogonadotropic hypogonadism (Kallmann Syndrome) alters GnRH neuronal migration. *Endocrinology*. 2016;157:1956-1966. Epub 2016 Mar 25

8. Howard SR, Guasti L, Ruiz-Babot G, Mancini A, David A, Storr HL, Metherell LA, Sternberg MJ, Cabrera CP, Warren HR, Barnes MR, Quinton R, de Roux N, Young J, Guiochon-Mantel A, Wehkalampi K, André V, Gothilf Y, Cariboni A, Dunkel L. IGSF10 mutations dysregulate gonadotropin-releasing hormone neuronal migration resulting in delayed puberty. *EMBO Mol Med*. 2016;8:626-642.
9. Palmert MR, Dunkel L. Clinical practice. Delayed puberty. *N Engl J Med*. 2012;366:443-453.
10. Sedlmeyer IL, Palmert MR. Delayed puberty: analysis of a large case series from an academic center. *J Clin Endocrinol Metab*. 2002;87:1613-1620.
11. Zhu J, Choa RE, Guo MH, Plummer L, Buck C, Palmert MR, Hirschhorn JN, Seminara SB, Chan YM. A shared genetic basis for self-limited delayed puberty and idiopathic hypogonadotropic hypogonadism. *J Clin Endocrinol Metab*. 2015;100:E646-E654. Epub 2015 Jan 30
12. Raivio T, Falardeau J, Dwyer A, Quinton R, Hayes FJ, Hughes VA, Cole LW, Pearce SH, Lee H, Boepple P, Crowley WF Jr, Pitteloud N. Reversal of idiopathic hypogonadotropic hypogonadism. *N Engl J Med*. 2007;357:863-873.
13. Sidhoum VF, Chan YM, Lippincott MF, Balasubramanian R, Quinton R, Plummer L, Dwyer A, Pitteloud N, Hayes FJ, Hall JE, Martin KA, Boepple PA, Seminara SB. Reversal and relapse of hypogonadotropic hypogonadism: resilience and fragility of the reproductive neuroendocrine system. *J Clin Endocrinol Metab*. 2014;99:861-870. Epub 2013 Jan 1
14. Lippincott MF, Schafer EC, Hindman AA, He W, Brauner R, Delaney A, Grinspon R, Hall JE, Hirschhorn JN, McElreavey K, Palmert MR, Rey R, Seminara SB, Salem RM, Chan YM; Delayed Puberty Genetics Consortium. Contributions of common genetic variants to constitutional delay of puberty and idiopathic hypogonadotropic hypogonadism. *J Clin Endocrinol Metab*. 2024;110:e61-e67.
15. Rezende RC, He W, Kaisinger LR, Lerario AM, Schafer EC, Kentistou KA, Barroso PS, Andrade NLM, Dantas NCB, Costa EMF, Cellin LP, P S Quedas E, Seminara SB, Rey RA, Grinspon RP, Meriq V, Ong KK, Latronico AC, Perry JRB, Howard SR, Chan YM, Jorge AAL; Delayed Puberty Genetics Consortium. Deleterious variants in intolerant genes reveal new candidates for self-limited delayed puberty. *Eur J Endocrinol*. 2025;192:481-490.
16. Duckett K, Williamson A, Kincaid JWR, Rainbow K, Corbin LJ, Martin HC, Eberhardt RY, Huang QQ, Hurler ME, He W, Brauner R, Delaney A, Dunkel L, Grinspon RP, Hall JE, Hirschhorn JN, Howard SR, Latronico AC, Jorge AAL, McElreavey K, Mericq V, Merino PM, Palmert MR, Plummer L, Rey RA, Rezende RC, Seminara SB, Salnikow K, Banerjee I, Lam BYH, Perry JRB, Timpson NJ, Clayton P, Chan YM, Ong KK, O'Rahilly S. Prevalence of deleterious variants in MC3R in patients with constitutional delay of growth and puberty. *J Clin Endocrinol Metab*. 2023;108:e1580-e1587.
17. Howard SR, Dunkel L. The Genetic Basis of Delayed Puberty. *Neuroendocrinology*. 2018;106:283-291. Epub 2017 Sep 18
18. Quaynor SD, Kim HG, Cappello EM, Williams T, Chorich LP, Bick DP, Sherins RJ, Layman LC. The prevalence of digenic mutations in patients with normosmic hypogonadotropic hypogonadism and Kallmann syndrome. *Fertil Steril*. 2011;96:1424-1430. Epub 2011 Oct 28
19. Pitteloud N, Quinton R, Pearce S, Raivio T, Acierno J, Dwyer A, Plummer L, Hughes V, Seminara S, Cheng YZ, Li WP, Maccoll G, Eliseenkova AV, Olsen SK, Ibrahim OA, Hayes FJ, Boepple P, Hall JE, Bouloux P, Mohammadi M, Crowley W. Digenic mutations account for variable phenotypes in idiopathic hypogonadotropic hypogonadism. *J Clin Invest*. 2007;117:457-463. Epub 2007 Jan 18
20. Sykiotis GP, Plummer L, Hughes VA, Au M, Durrani S, Nayak-Young S, Dwyer AA, Quinton R, Hall JE, Gusella JF, Seminara SB, Crowley WF Jr, Pitteloud N. Oligogenic basis of isolated gonadotropin-releasing hormone deficiency. *Proc Natl Acad Sci U S A*. 2010;107:15140-15144. Epub 2010 Aug 9
21. Boehm U, Bouloux PM, Dattani MT, de Roux N, Dodé C, Dunkel L, Dwyer AA, Giacobini P, Hardelin JP, Juul A, Maghnie M, Pitteloud N, Prevot V, Raivio T, Tena-Sempere M, Quinton R, Young J. Expert consensus document: European Consensus Statement on congenital hypogonadotropic hypogonadism--pathogenesis, diagnosis and treatment. *Nat Rev Endocrinol*. 2015;11:547-564. Epub 2015 Jul 21
22. Chong JX, Buckingham KJ, Jhangiani SN, Boehm C, Sobreira N, Smith JD, Harrell TM, McMillin MJ, Wiszniewski W, Gambin T, Coban Akdemir ZH, Doheny K, Scott AF, Avramopoulos D, Chakravarti A, Hoover-Fong J, Mathews D, Witmer PD, Ling H, Hetrick K, Watkins L, Patterson KE, Reinier F, Blue E, Muzny D, Kircher M, Bilguvar K, López-Giráldez F, Sutton VR, Tabor HK, Leal SM, Gunel M, Mane S, Gibbs RA, Boerwinkle E, Hamosh A, Shendure J, Lupski JR, Lifton RP, Valle D, Nickerson DA; Centers for Mendelian Genomics; Bamshad MJ. The genetic basis of mendelian phenotypes: discoveries, challenges, and opportunities. *Am J Hum Genet*. 2015;97:199-215. Epub 2015 Jul 9
23. Al Sayed Y, Howard SR. Panel testing for the molecular genetic diagnosis of congenital hypogonadotropic hypogonadism - a clinical perspective. *Eur J Hum Genet*. 2023;31:387-394. Epub 2022 Dec 15
24. Gürbüz F, Kotan LD, Mengen E, Şıklar Z, Berberoğlu M, Dökmetaş S, Kılıçlı MF, Güven A, Kirel B, Saka N, Poyrazoğlu Ş, Cesur Y, Doğan M, Özen S, Özbek MN, Demirebilek H, Kekil MB, Temiz F, Önelli Mungan N, Yüksel B, Topaloğlu AK. Distribution of gene mutations associated with familial normosmic idiopathic hypogonadotropic hypogonadism. *J Clin Res Pediatr Endocrinol*. 2012;4:121-126. Epub 2012 Jul 5
25. Quinton R, Duke VM, de Zoysa PA, Platts AD, Valentine A, Kendall B, Pickman S, Kirk JM, Besser GM, Jacobs HS, Bouloux PM. The neuroradiology of Kallmann's syndrome: a genotypic and phenotypic analysis. *J Clin Endocrinol Metab*. 1996;81:3010-3017.
26. Seminara SB, Hayes FJ, Crowley WF Jr. Gonadotropin-releasing hormone deficiency in the human (idiopathic hypogonadotropic hypogonadism and Kallmann's syndrome): pathophysiological and genetic considerations. *Endocr Rev*. 1998;19:521-539.
27. Nachtigall LB, Boepple PA, Pralong FP, Crowley WF Jr. Adult-onset idiopathic hypogonadotropic hypogonadism--a treatable form of male infertility. *N Engl J Med*. 1997;336:410-415.
28. Costa-Barbosa FA, Balasubramanian R, Keefe KW, Shaw ND, Al-Tassan N, Plummer L, Dwyer AA, Buck CL, Choi JH, Seminara SB, Quinton R, Monies D, Meyer B, Hall JE, Pitteloud N, Crowley WF Jr. Prioritizing genetic testing in patients with Kallmann syndrome using clinical phenotypes. *J Clin Endocrinol Metab*. 2013;98:E943-E953. Epub 2013 Mar 26
29. Dodé C, Levilliers J, Dupont JM, De Paepe A, Le Dü N, Soussi-Yanicostas N, Coimbra RS, Delmaghani S, Compain-Nouaille S, Baverel F, Pêcheux C, Le Tessier D, Cruaud C, Delpech M, Speleman F, Vermeulen S, Amalfitano A, Bachelot Y, Bouchard P, Cabrol S, Carel JC, Deleamarre-van de Waal H, Goulet-Salmon B, Pettit ML, Richard O, Sanchez-Franco F, Saura R, Young J, Kottler J, Hardelin JP. Loss-of-function mutations in FGFR1 cause autosomal dominant Kallmann syndrome. *Nat Genet*. 2003;33:463-465. Epub 2003 Mar 10
30. Hardelin JP, Dodé C. The complex genetics of Kallmann syndrome: KAL1, FGFR1, FGF8, PROKR2, PROK2, et al. *Sex Dev*. 2008;2:181-193. Epub 2008 Nov 5
31. Legouis R, Hardelin JP, Levilliers J, Clavier JM, Compain S, Wunderle V, Millasseau P, Le Paslier D, Cohen D, Caterina D, et al. The candidate gene for the X-linked Kallmann syndrome encodes a protein related to adhesion molecules. *Cell*. 1991;67:423-435.
32. de Castro F, Seal R, Maggi R; Group of HGNC consultants for KAL1 nomenclature. ANOS1: a unified nomenclature for Kallmann syndrome 1 gene (KAL1) and anosmin-1. *Brief Funct Genomics*. 2017;16:205-210.
33. Pedersen-White JR, Chorich LP, Bick DP, Sherins RJ, Layman LC. The prevalence of intragenic deletions in patients with idiopathic hypogonadotropic hypogonadism and Kallmann syndrome. *Mol Hum Reprod*. 2008;14:367-370. Epub 2008 May 7

34. Oliveira LM, Seminara SB, Beranova M, Hayes FJ, Valkenburgh SB, Schipani E, Costa EM, Latronico AC, Crowley WF Jr, Vallejo M. The importance of autosomal genes in Kallmann syndrome: genotype-phenotype correlations and neuroendocrine characteristics. *J Clin Endocrinol Metab.* 2001;86:1532-1538.
35. Franco B, Guioli S, Pragliola A, Incerti B, Bardoni B, Tonlorenzi R, Carozzo R, Maestrini E, Pieretti M, Taillon-Miller P, Brown CJ, Willard HF, Lawrence C, Graziella Persico M, Camerino G, Ballabio A. A gene deleted in Kallmann's syndrome shares homology with neural cell adhesion and axonal path-finding molecules. *Nature.* 1991;353:529-536.
36. Esteban PF, Murcia-Belmonte V, García-González D, de Castro F. The cysteine-rich region and the whey acidic protein domain are essential for anosmin-1 biological functions. *J Neurochem.* 2013;124:708-720. Epub 2012 Dec 21
37. Balasubramanian R, Crowley WF, Jr. Isolated Gonadotropin-Releasing Hormone (GnRH) Deficiency. In: Adam MP, Feldman J, Mirzaa GM, Pagon RA, Wallace SE, Amemiya A (eds). *GeneReviews*(®). Seattle (WA),1993.
38. Schwanzel-Fukuda M, Bick D, Pfaff DW. Luteinizing hormone-releasing hormone (LHRH)-expressing cells do not migrate normally in an inherited hypogonadal (Kallmann) syndrome. *Brain Res Mol Brain Res.* 1989;6:311-326.
39. Gonçalves CI, Fonseca F, Borges T, Cunha F, Lemos MC. Expanding the genetic spectrum of ANOS1 mutations in patients with congenital hypogonadotropic hypogonadism. *Hum Reprod.* 2017;32:704-711.
40. Nie M, Xu H, Chen R, Mao J, Wang X, Xiong S, Zheng J, Yu B, Cui M, Ma W, Huang Q, Zhang H, Wu X. Analysis of genetic and clinical characteristics of a Chinese Kallmann syndrome cohort with ANOS1 mutations. *Eur J Endocrinol.* 2017;177:389-398. Epub 2017 Aug 5
41. Tsai PS, Gill JC. Mechanisms of disease: Insights into X-linked and autosomal-dominant Kallmann syndrome. *Nat Clin Pract Endocrinol Metab.* 2006;2:160-171.
42. Vermeulen S, Messiaen L, Scheir P, De Bie S, Speleman F, De Paep A. Kallmann syndrome in a patient with congenital spherocytosis and an interstitial 8p11.2 deletion. *Am J Med Genet.* 2002;108:315-318.
43. Trarbach EB, Costa EM, Versiani B, de Castro M, Baptista MT, Garmes HM, de Mendonca BB, Latronico AC. Novel fibroblast growth factor receptor 1 mutations in patients with congenital hypogonadotropic hypogonadism with and without anosmia. *J Clin Endocrinol Metab.* 2006;91:4006-4012. Epub 2006 Aug 1 Erratum in *J Clin Endocrinol Metab.* 2008;93:2013.
44. Xu N, Qin Y, Reindollar RH, Tho SP, McDonough PG, Layman LC. A mutation in the fibroblast growth factor receptor 1 gene causes fully penetrant normosmic isolated hypogonadotropic hypogonadism. *J Clin Endocrinol Metab.* 2007;92:1155-1158. Epub 2007 Jan 2
45. Pitteloud N, Meysing A, Quinton R, Acierno JS Jr, Dwyer AA, Plummer L, Fliers E, Boepple P, Hayes F, Seminara S, Hughes VA, Ma J, Bouloux P, Mohammadi M, Crowley WF Jr. Mutations in fibroblast growth factor receptor 1 cause Kallmann syndrome with a wide spectrum of reproductive phenotypes. *Mol Cell Endocrinol.* 2006;254-255:60-69. Epub 2006 Jun 9
46. Raivio T, Sidis Y, Plummer L, Chen H, Ma J, Mukherjee A, Jacobson-Dickman E, Quinton R, Van Vliet G, Lavoie H, Hughes VA, Dwyer A, Hayes FJ, Xu S, Sparks S, Kaiser UB, Mohammadi M, Pitteloud N. Impaired fibroblast growth factor receptor 1 signaling as a cause of normosmic idiopathic hypogonadotropic hypogonadism. *J Clin Endocrinol Metab.* 2009;94:4380-4390. Epub 2009 Oct 9
47. Akkuş G, Kotan LD, Durmaz E, Mengen E, Turan İ, Ulubay A, Gürbüz F, Yüksel B, Tetiker T, Topaloglu AK. Hypogonadotropic Hypogonadism due to Novel FGFR1 Mutations. *J Clin Res Pediatr Endocrinol.* 2017;9:95-100. Epub 2016 Dec 23
48. Topaloglu AK, Kotan LD. Genetics of Hypogonadotropic hypogonadism. *Endocr Dev.* 2016;29:36-49. Epub 2015 Dec 17
49. Pitteloud N, Acierno JS Jr, Meysing AU, Dwyer AA, Hayes FJ, Crowley WF Jr. Reversible kallmann syndrome, delayed puberty, and isolated anosmia occurring in a single family with a mutation in the fibroblast growth factor receptor 1 gene. *J Clin Endocrinol Metab.* 2005;90:1317-1322. Epub 2004 Dec 21
50. Pitteloud N, Acierno JS Jr, Meysing A, Eliseenkova AV, Ma J, Ibrahim OA, Metzger DL, Hayes FJ, Dwyer AA, Hughes VA, Yialamas M, Hall JE, Grant E, Mohammadi M, Crowley WF Jr. Mutations in fibroblast growth factor receptor 1 cause both Kallmann syndrome and normosmic idiopathic hypogonadotropic hypogonadism. *Proc Natl Acad Sci U S A.* 2006;103:6281-6286. Epub 2006 Apr 10
51. Sato N, Katsumata N, Kagami M, Hasegawa T, Hori N, Kawakita S, Minowada S, Shimotsuka A, Shishiba Y, Yokozawa M, Yasuda T, Nagasaki K, Hasegawa D, Hasegawa Y, Tachibana K, Naiki Y, Horikawa R, Tanaka T, Ogata T. Clinical assessment and mutation analysis of Kallmann syndrome 1 (KAL1) and fibroblast growth factor receptor 1 (FGFR1, or KAL2) in five families and 18 sporadic patients. *J Clin Endocrinol Metab.* 2004;89:1079-1088.
52. Falardeau J, Chung WC, Beenken A, Raivio T, Plummer L, Sidis Y, Jacobson-Dickman EE, Eliseenkova AV, Ma J, Dwyer A, Quinton R, Na S, Hall JE, Huot C, Alois N, Pearce SH, Cole LW, Hughes V, Mohammadi M, Tsai P, Pitteloud N. Decreased FGF8 signaling causes deficiency of gonadotropin-releasing hormone in humans and mice. *J Clin Invest.* 2008;118:2822-2831.
53. Miraoui H, Dwyer AA, Sykiotis GP, Plummer L, Chung W, Feng B, Beenken A, Clarke J, Pers TH, Dworzynski P, Keefe K, Niedziela M, Raivio T, Crowley WF Jr, Seminara SB, Quinton R, Hughes VA, Kumanov P, Young J, Yialamas MA, Hall JE, Van Vliet G, Chanoine JP, Rubenstein J, Mohammadi M, Tsai PS, Sidis Y, Lage K, Pitteloud N. Mutations in FGF17, IL17RD, DUSP6, SPRY4, and FLRT3 are identified in individuals with congenital hypogonadotropic hypogonadism. *Am J Hum Genet.* 2013;92:725-743.
54. Trarbach EB, Abreu AP, Silveira LF, Garmes HM, Baptista MT, Teles MG, Costa EM, Mohammadi M, Pitteloud N, Mendonca BB, Latronico AC. Nonsense mutations in FGF8 gene causing different degrees of human gonadotropin-releasing deficiency. *J Clin Endocrinol Metab.* 2010;95:3491-3496. Epub 2010 May 12 Erratum in: *J Clin Endocrinol Metab.* 2011;96:2624.
55. Chung WC, Matthews TA, Tata BK, Tsai PS. Compound deficiencies in multiple fibroblast growth factor signalling components differentially impact the murine gonadotrophin-releasing hormone system. *J Neuroendocrinol.* 2010;22:944-950. Epub 2010 Jun 9
56. Xu C, Messina A, Somm E, Miraoui H, Kinnunen T, Acierno J Jr, Niederländer NJ, Bouilly J, Dwyer AA, Sidis Y, Cassatella D, Sykiotis GP, Quinton R, De Geyter C, Dirlwanger M, Schwitzgebel V, Cole TR, Toogood AA, Kirk JM, Plummer L, Albrecht U, Crowley WF Jr, Mohammadi M, Tena-Sempere M, Prevot V, Pitteloud N. *KLB*, encoding β -Klotho, is mutated in patients with congenital hypogonadotropic hypogonadism. *EMBO Mol Med.* 2017;9:1379-1397.
57. González-Martínez D, Kim SH, Hu Y, Guimond S, Schofield J, Winyard P, Vannelli GB, Turnbull J, Bouloux PM. Anosmin-1 modulates fibroblast growth factor receptor 1 signaling in human gonadotropin-releasing hormone olfactory neuroblasts through a heparan sulfate-dependent mechanism. *J Neurosci.* 2004;24:10384-10392.
58. Tornberg J, Sykiotis GP, Keefe K, Plummer L, Hoang X, Hall JE, Quinton R, Seminara SB, Hughes V, Van Vliet G, Van Uum S, Crowley WF, Habuchi H, Kimata K, Pitteloud N, Bülow HE. Heparan sulfate 6-O-sulfotransferase 1, a gene involved in extracellular sugar modifications, is mutated in patients with idiopathic hypogonadotropic hypogonadism. *Proc Natl Acad Sci U S A.* 2011;108:11524-11529. Epub 2011 Jun 23
59. Cheng MY, Leslie FM, Zhou QY. Expression of prokineticins and their receptors in the adult mouse brain. *J Comp Neurol.* 2006;498:796-809.
60. Ng KL, Li JD, Cheng MY, Leslie FM, Lee AG, Zhou QY. Dependence of olfactory bulb neurogenesis on prokineticin 2 signaling. *Science.* 2005;308:1923-1927.

61. Pitteloud N, Zhang C, Pignatelli D, Li JD, Raivio T, Cole LW, Plummer L, Jacobson-Dickman EE, Mellon PL, Zhou QY, Crowley WF Jr. Loss-of-function mutation in the prokineticin 2 gene causes Kallmann syndrome and normosmic idiopathic hypogonadotropic hypogonadism. *Proc Natl Acad Sci U S A*. 2007;104:17447-17452. Epub 2007 Oct 24
62. Matsumoto S, Yamazaki C, Masumoto KH, Nagano M, Naito M, Soga T, Hiyama H, Matsumoto M, Takasaki J, Kamohara M, Matsuo A, Ishii H, Kobori M, Katoh M, Matsushime H, Furuichi K, Shigeyoshi Y. Abnormal development of the olfactory bulb and reproductive system in mice lacking prokineticin receptor PKR2. *Proc Natl Acad Sci U S A*. 2006;103:4140-4145. Epub 2006 Mar 2
63. Dodé C, Teixeira L, Levilliers J, Fouveaut C, Bouchard P, Kottler ML, Lespinasse J, Lienhardt-Roussie A, Mathieu M, Moerman A, Morgan G, Murat A, Toublanc JE, Wolczynski S, Delpech M, Petit C, Young J, Hardelin JP. Kallmann syndrome: mutations in the genes encoding prokineticin-2 and prokineticin receptor-2. *PLoS Genet*. 2006;2:e175. Epub 2006 Sep 1
64. Abreu AP, Trarbach EB, de Castro M, Frade Costa EM, Versiani B, Matias Baptista MT, Garmes HM, Mendonca BB, Latronico AC. Loss-of-function mutations in the genes encoding prokineticin-2 or prokineticin receptor-2 cause autosomal recessive Kallmann syndrome. *J Clin Endocrinol Metab*. 2008;93:4113-4118. Epub 2008 Aug 5
65. Cole LW, Sidis Y, Zhang C, Quinton R, Plummer L, Pignatelli D, Hughes VA, Dwyer AA, Raivio T, Hayes FJ, Seminara SB, Huot C, Alos N, Speiser P, Takeshita A, Van Vliet G, Pearce S, Crowley WF Jr, Zhou QY, Pitteloud N. Mutations in prokineticin 2 and prokineticin receptor 2 genes in human gonadotrophin-releasing hormone deficiency: molecular genetics and clinical spectrum. *J Clin Endocrinol Metab*. 2008;93:3551-3559. Epub 2008 Jun 17
66. Stamou MI, Chiu CJ, Jadhav SV, Salnikov KB, Plummer L, Seminara SB, Balasubramanian R. Expanding the phenotypic spectrum of PROK2/PROKR2: a recall-by-genotype study. *Hum Genet*. 2025;144:761-773. Epub 2025 Jun 11
67. Vissers LE, van Ravenswaaij CM, Admiraal R, Hurst JA, de Vries BB, Janssen IM, van der Vliet WA, Huys EH, de Jong PJ, Hamel BC, Schoenmakers EF, Brunner HG, Veltman JA, van Kessel AG. Mutations in a new member of the chromodomain gene family cause CHARGE syndrome. *Nat Genet*. 2004;36:955-957. Epub 2004 Aug 8
68. Layman WS, Hurd EA, Martin DM. Reproductive dysfunction and decreased GnRH neurogenesis in a mouse model of CHARGE syndrome. *Hum Mol Genet*. 2011;20:3138-3150. Epub 2011 May 19
69. Pinto G, Abadie V, Mesnage R, Blustajn J, Cabrol S, Amiel J, Hertz-Pannier L, Bertrand AM, Lyonnet S, Rappaport R, Netchine I. CHARGE syndrome includes hypogonadotropic hypogonadism and abnormal olfactory bulb development. *J Clin Endocrinol Metab*. 2005;90:5621-5626. Epub 2005 Jul 19
70. Marcos S, Sarfati J, Leroy C, Fouveaut C, Parent P, Metz C, Wolczynski S, Gérard M, Bieth E, Kurtz F, Verier-Mine O, Perrin L, Archambeaud F, Cabrol S, Rodien P, Hove H, Prescott T, Lacombe D, Christin-Maitre S, Touraine P, Hieronimus S, Dewailly D, Young J, Pugeat M, Hardelin JP, Dodé C. The prevalence of CHD7 missense versus truncating mutations is higher in patients with Kallmann syndrome than in typical CHARGE patients. *J Clin Endocrinol Metab*. 2014;99:E2138-E2143. Epub 2014 Jul 31 Erratum in: *J Clin Endocrinol Metab*. 2015;100:317.
71. Balasubramanian R, Choi JH, Francescato L, Willer J, Horton ER, Asimacopoulos EP, Stankovic KM, Plummer L, Buck CL, Quinton R, Nebesio TD, Mericq V, Merino PM, Meyer BF, Monies D, Gusella JF, Al Tassan N, Katsanis N, Crowley WF Jr. Functionally compromised CHD7 alleles in patients with isolated GnRH deficiency. *Proc Natl Acad Sci U S A*. 2014;111:17953-17958. Epub 2014 Dec 3
72. Kim YJ, Osborn DP, Lee JY, Araki M, Araki K, Mohun T, Käsäkoski J, Brandstäck N, Kim HT, Miralles F, Kim CH, Brown NA, Kim HG, Martínez-Barbera JP, Ataliotis P, Raivio T, Layman LC, Kim SH. WDR11-mediated Hedgehog signalling defects underlie a new ciliopathy related to Kallmann syndrome. *EMBO Rep*. 2018;19:269-289. Epub 2017 Dec 20
73. Kim HG, Ahn JW, Kurth I, Ullmann R, Kim HT, Kulharya A, Ha KS, Itokawa Y, Meliciani I, Wenzel W, Lee D, Rosenberger G, Ozata M, Bick DP, Sherins RJ, Nagase T, Tekin M, Kim SH, Kim CH, Ropers HH, Gusella JF, Kalscheuer V, Choi CY, Layman LC. WDR11, a WD protein that interacts with transcription factor EMX1, is mutated in idiopathic hypogonadotropic hypogonadism and Kallmann syndrome. *Am J Hum Genet*. 2010;87:465-479.
74. Turan I, Hutchins BI, Hacıhamdioglu B, Kotan LD, Gurbuz F, Ulubay A, Mengen E, Yuksel B, Wray S, Topaloglu AK. CCDC141 mutations in idiopathic hypogonadotropic hypogonadism. *J Clin Endocrinol Metab*. 2017;102:1816-1825.
75. Saengkaew T, Ruiz-Babot G, David A, Mancini A, Mariniello K, Cabrera CP, Barnes MR, Dunkel L, Guasti L, Howard SR. Whole exome sequencing identifies deleterious rare variants in CCDC141 in familial self-limited delayed puberty. *NPJ Genom Med*. 2021;6:107.
76. Eckler MJ, McKenna WL, Taghvaei S, McConnell SK, Chen B. Fezf1 and Fezf2 are required for olfactory development and sensory neuron identity. *J Comp Neurol*. 2011;519:1829-1846.
77. Watanabe Y, Inoue K, Okuyama-Yamamoto A, Nakai N, Nakatani J, Nibu K, Sato N, Iiboshi Y, Yusa K, Kondoh G, Takeda J, Terashima T, Takumi T. Fezf1 is required for penetration of the basal lamina by olfactory axons to promote olfactory development. *J Comp Neurol*. 2009;515:565-584.
78. Kotan LD, Hutchins BI, Ozkan Y, Demirel F, Stoner H, Cheng PJ, Esen I, Gurbuz F, Bicakci YK, Mengen E, Yuksel B, Wray S, Topaloglu AK. Mutations in FEZF1 cause Kallmann syndrome. *Am J Hum Genet*. 2014;95:326-331.
79. Giacobini P. Shaping the reproductive system: role of semaphorins in gonadotropin-releasing hormone development and function. *Neuroendocrinology*. 2015;102:200-215. Epub 2015 May 7
80. Oleari R, André V, Lettieri A, Tahir S, Roth L, Paganoni A, Eberini I, Parravicini C, Scagliotti V, Cotellessa L, Bedogni F, De Martini LB, Corridor MV, Gulli S, Augustin HG, Gaston-Massuet C, Hussain K, Cariboni A. A novel SEMA3G mutation in two siblings affected by syndromic GnRH deficiency. *Neuroendocrinology*. 2021;111:421-441. Epub 2020 May 4
81. Cariboni A, André V, Chauvet S, Cassatella D, Davidson K, Caramello A, Fantin A, Bouloux P, Mann F, Ruhrberg C. Dysfunctional SEMA3E signaling underlies gonadotropin-releasing hormone neuron deficiency in Kallmann syndrome. *J Clin Invest*. 2015;125:2413-2428. Epub 2015 May 18
82. Hanchate NK, Giacobini P, Lhuillier P, Parkash J, Espy C, Fouveaut C, Leroy C, Baron S, Campagne C, Vanacker C, Collier F, Cruaud C, Meyer V, García-Piñero A, Dewailly D, Cortet-Rudelli C, Gersak K, Metz C, Chabrier G, Pugeat M, Young J, Hardelin JP, Prevot V, Dodé C. SEMA3A, a gene involved in axonal pathfinding, is mutated in patients with Kallmann syndrome. *PLoS Genet*. 2012;8:e1002896. Epub 2012 Aug 23
83. Janssen BJ, Malinauskas T, Weir GA, Cader MZ, Siebold C, Jones EY. Neuropilins lock secreted semaphorins onto plexins in a ternary signaling complex. *Nat Struct Mol Biol*. 2012;19:1293-1299. Epub 2012 Oct 28
84. Marcos S, Monnier C, Rovira X, Fouveaut C, Pitteloud N, Ango F, Dodé C, Hardelin JP. Defective signaling through plexin-A1 compromises the development of the peripheral olfactory system and neuroendocrine reproductive axis in mice. *Hum Mol Genet*. 2017;26:2006-2017.
85. Kotan LD, Ternier G, Cakir AD, Emekci HC, Turan I, Delpouve G, Kardelen AD, Ozcabi B, Isik E, Mengen E, Cakir EDP, Yuksel A, Agladioglu SY, Dilek SO, Evliyaoglu O, Darendeliler F, Gurbuz F, Akkus G, Yuksel B, Giacobini P, Topaloglu AK. Loss-of-function variants in SEMA3F and PLXNA3 encoding semaphorin-3F and its receptor plexin-A3 respectively cause idiopathic hypogonadotropic hypogonadism. *Genet Med*. 2021;23:1008-1016. Epub 2021 Jan 25

86. Limoni G, Niquille M. Semaphorins and Plexins in central nervous system patterning: the key to it all? *Curr Opin Neurobiol.* 2021;66:224-232. Epub 2021 Jan 26
87. Worzfeld T, Offermanns S. Semaphorins and plexins as therapeutic targets. *Nat Rev Drug Discov.* 2014;13:603-621.
88. Kotan LD, Isik E, Turan I, Mengen E, Akkus G, Tastan M, Gurbuz F, Yuksel B, Topaloglu AK. Prevalence and associated phenotypes of PLXNA1 variants in normosmic and anosmic idiopathic hypogonadotropic hypogonadism. *Clin Genet.* 2019;95:320-324. Epub 2018 Dec 26
89. Welch BA, Cho HJ, Ucakurk SA, Farmer SM, Cetinkaya S, Abaci A, Akkus G, Simsek E, Kotan LD, Turan I, Gurbuz F, Yuksel B, Wray S, Topaloglu AK. PLXNB1 mutations in the etiology of idiopathic hypogonadotropic hypogonadism. *J Neuroendocrinol.* 2022;34:e13103. Epub 2022 Feb 16
90. Shaw ND, Brand H, Kupchinsky ZA, Bengani H, Plummer L, Jones TI, Erdin S, Williamson KA, Rainger J, Stortchevoi A, Samocha K, Currall BB, Dunican DS, Collins RL, Willer JR, Lek A, Lek M, Nassan M, Pereira S, Kammin T, Lucente D, Silva A, Seabra CM, Chiang C, An Y, Ansari M, Rainger JK, Joss S, Smith JC, Lippincott MF, Singh SS, Patel N, Jing JW, Law JR, Ferraro N, Verloes A, Rauch A, Steindl K, Zweier M, Scheer I, Sato D, Okamoto N, Jacobsen C, Tryggstad J, Chernauek S, Schimmenti LA, Brasseur B, Cesaretti C, García-Ortiz JE, Buitrago TP, Silva OP, Hoffman JD, Mühlbauer W, Ruprecht KW, Loeys BL, Shino M, Kaindl AM, Cho CH, Morton CC, Meehan RR, van Heyningen V, Liao EC, Balasubramanian R, Hall JE, Seminara SB, Macarthur D, Moore SA, Yoshiura KI, Gusella JF, Marsh JA, Graham JM Jr, Lin AE, Katsanis N, Jones PL, Crowley WF Jr, Davis EE, FitzPatrick DR, Talkowski ME. SMCHD1 mutations associated with a rare muscular dystrophy can also cause isolated arhinia and Bosma arhinia microphthalmia syndrome. *Nat Genet.* 2017;49:238-248. Epub 2017 Jan 9 Erratum in: *Nat Genet.* 2017;49:969.
91. Pingault V, Bodereau V, Baral V, Marcos S, Watanabe Y, Chaoui A, Fouveaut C, Leroy C, Vêrier-Mine O, Francannet C, Dupin-Deguine D, Archambeaud F, Kurtz FJ, Young J, Bertherat J, Marlin S, Goossens M, Hardelin JP, Dodé C, Bondurand N. Loss-of-function mutations in SOX10 cause Kallmann syndrome with deafness. *Am J Hum Genet.* 2013;92:707-724.
92. Rojas RA, Kutateladze AA, Plummer L, Stamou M, Keefe DL Jr, Salnikov KB, Delaney A, Hall JE, Sadreyev R, Ji F, Fliers E, Gambosova K, Quinton R, Merino PM, Mericq V, Seminara SB, Crowley WF Jr, Balasubramanian R. Phenotypic continuum between Waardenburg syndrome and idiopathic hypogonadotropic hypogonadism in humans with SOX10 variants. *Genet Med.* 2021;23:629-636. Epub 2021 Jan 13
93. Cassin J, Stamou MI, Keefe KW, Sung KE, Bojo CC, Tonsfeldt KJ, Rojas RA, Ferreira Lopes V, Plummer L, Salnikov KB, Keefe DL Jr, Ozata M, Genel M, Georgopoulos NA, Hall JE, Crowley WF Jr, Seminara SB, Mellon PL, Balasubramanian R. Heterozygous mutations in SOX2 may cause idiopathic hypogonadotropic hypogonadism via dominant-negative mechanisms. *JCI Insight.* 2023;8:e164324.
94. Messina A, Pulli K, Santini S, Acierno J, Käsäkoski J, Cassatella D, Xu C, Casoni F, Malone SA, Ternier G, Conte D, Sidis Y, Tommiska J, Vaaralahti K, Dwyer A, Gothilf Y, Merlo GR, Santoni F, Niederländer NJ, Giacobini P, Raivio T, Pitteloud N. Neuron-derived neurotrophic factor is mutated in congenital hypogonadotropic hypogonadism. *Am J Hum Genet.* 2020;106:58-70. Epub 2019 Dec 26
95. Kotan LD, Yildiz M, Turan I, Celiloglu C, Yuksel B, Topaloglu AK. A novel homozygous nonsense NDNF variant in Kallmann syndrome. *Am J Med Genet A.* 2023;191:831-834. Epub 2022 Dec 1
96. Gowkielewicz M, Lipka A, Zdanowski W, Waśniewski T, Majewska M, Carlberg C. Anti-Müllerian hormone: biology and role in endocrinology and cancers. *Front Endocrinol (Lausanne).* 2024;15:1468364.
97. Malone SA, Papadakis GE, Messina A, Mimouni NEH, Trova S, Imbernon M, Allet C, Cimino I, Acierno J, Cassatella D, Xu C, Quinton R, Szinnai G, Pigny P, Alonso-Cotchico L, Masgrau L, Maréchal JD, Prevot V, Pitteloud N, Giacobini P. Defective AMH signaling disrupts GnRH neuron development and function and contributes to hypogonadotropic hypogonadism. *Elife.* 2019;8:e47198.
98. Salian-Mehta S, Xu M, Knox AJ, Plummer L, Slavov D, Taylor M, Bevers S, Hodges RS, Crowley WF Jr, Wierman ME. Functional consequences of AXL sequence variants in hypogonadotropic hypogonadism. *J Clin Endocrinol Metab.* 2014;99:1452-1460. Epub 2014 Jan 2
99. Schwarting GA, Kostek C, Bless EP, Ahmad N, Tobet SA. Deleted in colorectal cancer (DCC) regulates the migration of luteinizing hormone-releasing hormone neurons to the basal forebrain. *J Neurosci.* 2001;21:911-919.
100. Schwarting GA, Raitcheva D, Bless EP, Ackerman SL, Tobet S. Netrin 1-mediated chemoattraction regulates the migratory pathway of LHRH neurons. *Eur J Neurosci.* 2004;19:11-20.
101. Bouilly J, Messina A, Papadakis G, Cassatella D, Xu C, Acierno JS, Tata B, Sykiotis G, Santini S, Sidis Y, Elowe-Gruau E, Phan-Hug F, Hauschild M, Bouloux PM, Quinton R, Lang-Muritano M, Favre L, Marino L, Giacobini P, Dwyer AA, Niederländer NJ, Pitteloud N. DCC/NTN1 complex mutations in patients with congenital hypogonadotropic hypogonadism impair GnRH neuron development. *Hum Mol Genet.* 2018;27:359-372.
102. de Roux N, Young J, Misrahi M, Genet R, Chanson P, Schaison G, Milgrom E. A family with hypogonadotropic hypogonadism and mutations in the gonadotropin-releasing hormone receptor. *N Engl J Med.* 1997;337:1597-1602.
103. Layman LC, Cohen DP, Jin M, Xie J, Li Z, Reindollar RH, Bolbolan S, Bick DP, Sherins RR, Duck LW, Musgrove LC, Sellers JC, Neill JD. Mutations in gonadotropin-releasing hormone receptor gene cause hypogonadotropic hypogonadism. *Nat Genet.* 1998;18:14-15.
104. Gianetti E, Hall JE, Au MG, Kaiser UB, Quinton R, Stewart JA, Metzger DL, Pitteloud N, Mericq V, Merino PM, Levitsky LL, Izatt L, Lang-Muritano M, Fujimoto VY, Dluhy RG, Chase ML, Crowley WF Jr, Plummer L, Seminara SB. When genetic load does not correlate with phenotypic spectrum: lessons from the GnRH receptor (GNRHR). *J Clin Endocrinol Metab.* 2012;97:E1798-E1807. Epub 2012 Jun 28
105. Francou B, Paul C, Amazit L, Cartes A, Bouvattier C, Albarel F, Maiter D, Chanson P, Trabado S, Brailly-Tabard S, Brue T, Guiochon-Mantel A, Young J, Bouligand J. Prevalence of KISS1 Receptor mutations in a series of 603 patients with normosmic congenital hypogonadotropic hypogonadism and characterization of novel mutations: a single-centre study. *Hum Reprod.* 2016;31:1363-1374. Epub 2016 Apr 19
106. Seminara SB, Topaloglu AK. Review of human genetic and clinical studies directly relevant to GnRH signalling. *J Neuroendocrinol.* 2022;34:e13080. Epub 2021 Dec 31
107. Topaloglu AK, Lu ZL, Farooqi IS, Mungan NO, Yuksel B, O'Rahilly S, Millar RP. Molecular genetic analysis of normosmic hypogonadotropic hypogonadism in a Turkish population: identification and detailed functional characterization of a novel mutation in the gonadotropin-releasing hormone receptor gene. *Neuroendocrinology.* 2006;84:301-308. Epub 2006 Dec 19
108. Pitteloud N, Boepple PA, DeCruz S, Valkenburgh SB, Crowley WF Jr, Hayes FJ. The fertile eunuch variant of idiopathic hypogonadotropic hypogonadism: spontaneous reversal associated with a homozygous mutation in the gonadotropin-releasing hormone receptor. *J Clin Endocrinol Metab.* 2001;86:2470-2475.
109. Cattanach BM, Iddon CA, Charlton HM, Chiappa SA, Fink G. Gonadotrophin-releasing hormone deficiency in a mutant mouse with hypogonadism. *Nature.* 1977;269:338-340.
110. Mason AJ, Pitts SL, Nikolics K, Szonyi E, Wilcox JN, Seeburg PH, Stewart TA. The hypogonadal mouse: reproductive functions restored by gene therapy. *Science.* 1986;234:1372-1378.

111. Bouligand J, Gervan C, Tello JA, Brailly-Tabard S, Salenave S, Chanson P, Lombès M, Millar RP, Guiochon-Mantel A, Young J. Isolated familial hypogonadotropic hypogonadism and a GNRH1 mutation. *N Engl J Med*. 2009;360:2742-2748. Epub 2009 Jun 17
112. Chan YM, de Guillebon A, Lang-Muritano M, Plummer L, Cerrato F, Tsiaras S, Gaspert A, Lavoie HB, Wu CH, Crowley WF Jr, Amory JK, Pitteloud N, Seminara SB. GNRH1 mutations in patients with idiopathic hypogonadotropic hypogonadism. *Proc Natl Acad Sci U S A*. 2009;106:11703-11708. Epub 2009 Jun 30
113. Mengen E, Tunc S, Kotan LD, Nalbantoglu O, Demir K, Gurbuz F, Turan I, Seker G, Yuksel B, Topaloglu AK. Complete idiopathic hypogonadotropic hypogonadism due to homozygous GNRH1 mutations in the mutational hot spots in the region encoding the decapeptide. *Horm Res Paediatr*. 2016;85:107-111. Epub 2015 Nov 24
114. Seminara SB, Messenger S, Chatzidakis EE, Thresher RR, Acierno JS Jr, Shagoury JK, Bo-Abbas Y, Kuohung W, Schwinoof KM, Hendrick AG, Zahn D, Dixon J, Kaiser UB, Slaugenhaupt SA, Gusella JF, O'Rahilly S, Carlton MB, Crowley WF Jr, Aparicio SA, Colledge WH. The GPR54 gene as a regulator of puberty. *N Engl J Med*. 2003;349:1614-1627.
115. de Roux N, Genin E, Carel JC, Matsuda F, Chaussain JL, Milgrom E. Hypogonadotropic hypogonadism due to loss of function of the KISS1-derived peptide receptor GPR54. *Proc Natl Acad Sci U S A*. 2003;100:10972-10976. Epub 2003 Aug 27
116. Cerrato F, Shagoury J, Kralickova M, Dwyer A, Falardeau J, Ozata M, Van Vliet G, Bouloux P, Hall JE, Hayes FJ, Pitteloud N, Martin KA, Welt C, Seminara SB. Coding sequence analysis of GNRHR and GPR54 in patients with congenital and adult-onset forms of hypogonadotropic hypogonadism. *Eur J Endocrinol*. 2006;155(Suppl 1):3-10.
117. Topaloglu AK, Tello JA, Kotan LD, Ozbek MN, Yilmaz MB, Erdogan S, Gurbuz F, Temiz F, Millar RP, Yuksel B. Inactivating KISS1 mutation and hypogonadotropic hypogonadism. *N Engl J Med*. 2012;366:629-635.
118. Coutant R, Zaegel N, Illouz F, Levaillant L, Magnin F, Perrot J, Ait Tayeb AEK, Briet C, Kaiser UB, Bouligand J, Young J, Maione L. Reversal of congenital hypogonadotropic hypogonadism in two brothers carrying a homozygous KISS1 frameshift mutation. *Lancet Diabetes Endocrinol*. 2025;13:550-552. Epub 2025 Jun 3
119. Topaloglu AK, Reimann F, Guclu M, Yalin AS, Kotan LD, Porter KM, Serin A, Mungan NO, Cook JR, Imamoglu S, Akalin NS, Yuksel B, O'Rahilly S, Semple RK. TAC3 and TACR3 mutations in familial hypogonadotropic hypogonadism reveal a key role for Neurokinin B in the central control of reproduction. *Nat Genet*. 2009;41:354-358. Epub 2008 Dec 11
120. Topaloglu AK, Kotan LD. Molecular causes of hypogonadotropic hypogonadism. *Curr Opin Obstet Gynecol*. 2010;22:264-270.
121. Gianetti E, Tusset C, Noel SD, Au MG, Dwyer AA, Hughes VA, Abreu AP, Carroll J, Trarbach E, Silveira LF, Costa EM, de Mendonça BB, de Castro M, Lofrano A, Hall JE, Bolu E, Ozata M, Quinton R, Amory JK, Stewart SE, Arlt W, Cole TR, Crowley WF, Kaiser UB, Latronico AC, Seminara SB. TAC3/TACR3 mutations reveal preferential activation of gonadotropin-releasing hormone release by neurokinin B in neonatal life followed by reversal in adulthood. *J Clin Endocrinol Metab*. 2010;95:2857-2867. Epub 2010 Mar 23
122. Francou B, Bouligand J, Voican A, Amazit L, Trabado S, Fagart J, Meduri G, Brailly-Tabard S, Chanson P, Lecomte P, Guiochon-Mantel A, Young J. Normosmic congenital hypogonadotropic hypogonadism due to TAC3/TACR3 mutations: characterization of neuroendocrine phenotypes and novel mutations. *PLoS One*. 2011;6:e25614. Epub 2011 Oct 21
123. Vaaralahti K, Wehkalampi K, Tommiska J, Laitinen EM, Dunkel L, Raivio T. The role of gene defects underlying isolated hypogonadotropic hypogonadism in patients with constitutional delay of growth and puberty. *Fertil Steril*. 2011;95:2756-2758. Epub 2011 Feb 3
124. Young J, Bouligand J, Francou B, Raffin-Sanson ML, Gaillez S, Jeanpierre M, Grynberg M, Kamenicky P, Chanson P, Brailly-Tabard S, Guiochon-Mantel A. TAC3 and TACR3 defects cause hypothalamic congenital hypogonadotropic hypogonadism in humans. *J Clin Endocrinol Metab*. 2010;95:2287-2295. Epub 2010 Mar 1
125. Perry JR, Day F, Elks CE, Sulem P, Thompson DJ, Ferreira T, He C, Chasman DI, Esko T, Thorleifsson G, Albrecht E, Ang WQ, Corre T, Cousminer DL, Feenstra B, Franceschini N, Ganna A, Johnson AD, Kjellqvist S, Lunetta KL, McMahon G, Nolte IM, Paternoster L, Porcu E, Smith AV, Stolk L, Teumer A, Tšernikova N, Tikkanen E, Ulivi S, Wagner EK, Amin N, Bierut LJ, Byrne EM, Hottenga JJ, Koller DL, Mangino M, Pers TH, Yerges-Armstrong LM, Zhao JH, Andrusil IL, Anton-Culver H, Atsma F, Bandinelli S, Beckmann MW, Benitez J, Blomqvist C, Bojesen SE, Bolla MK, Bonanni B, Brauch H, Brenner H, Buring JE, Chang-Claude J, Chanock S, Chen J, Chenevix-Trench G, Collée JM, Couch FJ, Couper D, Coveillo AD, Cox A, Czene K, D'Adamo AP, Smith GD, De Vivo I, Demerath EW, Dennis J, Devilee P, Dieffenbach AK, Dunning AM, Eiriksdottir G, Eriksson JG, Fasching PA, Ferrucci L, Flesch-Janys D, Flyger H, Forud T, Franke L, Garcia ME, García-Closas M, Geller F, de Geus EE, Giles GG, Gudbjartsson DF, Gudnason V, Guénel P, Guo S, Hall P, Hamann U, Haring R, Hartman CA, Heath AC, Hofman A, Hoening MJ, Hopper JL, Hu FB, Hunter DJ, Karasik D, Kiel DP, Knight JA, Kosma VM, Kutalik Z, Lai S, Lambrechts D, Lindblom A, Mägi R, Magnusson PK, Mannermaa A, Martin NG, Masson G, McArdle PF, McArdle WL, Melbye M, Michailidou K, Mihailov E, Milani L, Milne RL, Nevanlinna H, Neven P, Nohr EA, Oldehinkel AJ, Oostra BA, Palotie A, Peacock M, Pedersen NL, Peterlongo P, Peto J, Pharoah PD, Postma DS, Pouta A, Pyrkäs K, Radice P, Ring S, Rivadeneira F, Robino A, Rose LM, Rudolph A, Salomaa V, Sanna S, Schlessinger D, Schmidt MK, Southey MC, Sovio U, Stampfer MJ, Stöckl D, Storniolo AM, Timpson NJ, Tyrer J, Visser JA, Vollenweider P, Völzke H, Waeber G, Waldenberger M, Wallaschowski H, Wang Q, Willemsen G, Winqvist R, Wolffenbuttel BH, Wright MJ; Australian Ovarian Cancer Study; GENICA Network; kConFab; LifeLines Cohort Study; InterAct Consortium; Early Growth Genetics (EGG) Consortium; Boomsma DI, Econs MJ, Khaw KT, Loos RJ, McCarthy MI, Montgomery GW, Rice JP, Streeten EA, Thorsteinsdottir U, van Duijn CM, Alizadeh BZ, Bergmann S, Boerwinkle E, Boyd HA, Crisponi L, Gasparini P, Gieger C, Harris TB, Ingelsson E, Järvelin MR, Kraft P, Lawlor D, Metspalu A, Pennell CE, Ridker PM, Snieder H, Sørensen TI, Spector TD, Strachan DP, Uitterlinden AG, Wareham NJ, Widen E, Zygmont M, Murray A, Easton DF, Stefansson K, Murabito JM, Ong KK. Parent-of-origin-specific allelic associations among 106 genomic loci for age at menarche. *Nature*. 2014;514:92-97. Epub 2014 Jul 23
126. Heger S, Mastronardi C, Dissen GA, Lomniczi A, Cabrera R, Roth CL, Jung H, Galimi F, Sippell W, Ojeda SR. Enhanced at puberty 1 (EAP1) is a new transcriptional regulator of the female neuroendocrine reproductive axis. *J Clin Invest*. 2007;117:2145-2154.
127. Mancini A, Howard SR, Cabrera CP, Barnes MR, David A, Wehkalampi K, Heger S, Lomniczi A, Guasti L, Ojeda SR, Dunkel L. EAP1 regulation of GnRH promoter activity is important for human pubertal timing. *Hum Mol Genet*. 2019;28:1357-1368.
128. Strobel A, Issad T, Camoin L, Ozata M, Strosberg AD. A leptin missense mutation associated with hypogonadism and morbid obesity. *Nat Genet*. 1998;18:213-215.
129. Farooqi IS, Wangenstein T, Collins S, Kimber W, Matarese G, Keogh JM, Lank E, Bottomley B, Lopez-Fernandez J, Ferraz-Amaro I, Dattani MT, Ercan O, Myhre AG, Retterstol L, Stanhope R, Edge JA, McKenzie S, Lessan N, Ghodsi M, De Rosa V, Perna F, Fontana S, Barroso I, Undlien DE, O'Rahilly S. Clinical and molecular genetic spectrum of congenital deficiency of the leptin receptor. *N Engl J Med*. 2007;356:237-247.
130. Farooqi IS, Jebb SA, Langmack G, Lawrence E, Cheetham CH, Prentice AM, Hughes IA, McCamish MA, O'Rahilly S. Effects of recombinant leptin therapy in a child with congenital leptin deficiency. *N Engl J Med*. 1999;341:879-884.
131. Iyer AK, McCabe ER. Molecular mechanisms of DAX1 action. *Mol Genet Metab*. 2004;83:60-73.

132. Achermann JC, Gu WX, Kotlar TJ, Meeks JJ, Sabacan LP, Seminara SB, Habiby RL, Hindmarsh PC, Bick DP, Sherins RJ, Crowley WF Jr, Layman LC, Jameson JL. Mutational analysis of DAX1 in patients with hypogonadotropic hypogonadism or pubertal delay. *J Clin Endocrinol Metab.* 1999;84:4497-4500.
133. Durmaz E, Turkkahraman D, Berdeli A, Atan M, Karaguzel G, Akcurin S, Bircan I. A novel DAX-1 mutation presented with precocious puberty and hypogonadotropic hypogonadism in different members of a large pedigree. *J Pediatr Endocrinol Metab.* 2013;26:551-555.
134. Darcan S, Goksen D, Ozen S, Ozkinay F, Durmaz B, Lalli E. Gonadotropin-dependent precocious puberty in a patient with X-linked adrenal hypoplasia congenita caused by a novel DAX-1 mutation. *Horm Res Paediatr.* 2011;75:153-156. Epub 2010 Oct 22
135. Loke KY, Poh LK, Lee WW, Lai PS. A case of X-linked adrenal hypoplasia congenita, central precocious puberty and absence of the DAX-1 gene: implications for pubertal regulation. *Horm Res.* 2009;71:298-304. Epub 2009 Apr 1
136. Rosenfeld MG, Glass CK. Coregulator codes of transcriptional regulation by nuclear receptors. *J Biol Chem.* 2001;276:36865-36868. Epub 2001 Jul 17
137. Sablin EP, Woods A, Krylova IN, Hwang P, Ingraham HA, Fletterick RJ. The structure of corepressor Dax-1 bound to its target nuclear receptor LHR-1. *Proc Natl Acad Sci U S A.* 2008;105:18390-18395. Epub 2008 Nov 17
138. Ramos-Pittol JM, Fernandes-Freitas I, Milona A, Manchishi SM, Rainbow K, Lam BYH, Tadross JA, Beucher A, Colledge WH, Cebola I, Murphy KG, Miguel-Aliaga I, Yeo GSH, Dhillo WS, Owen BM. Dax1 modulates ER α -dependent hypothalamic estrogen sensing in female mice. *Nat Commun.* 2023;14:3076.
139. Chooniedass-Kothari S, Emberley E, Hamedani MK, Troup S, Wang X, Czosnek A, Hube F, Mutawe M, Watson PH, Leygue E. The steroid receptor RNA activator is the first functional RNA encoding a protein. *FEBS Lett.* 2004;566:43-47.
140. Kelly VR, Xu B, Kuick R, Koenig RJ, Hammer GD. Dax1 up-regulates Oct4 expression in mouse embryonic stem cells via LHR-1 and SRA. *Mol Endocrinol.* 2010;24:2281-2291. Epub 2010 Oct 13
141. Kotan LD, Cooper C, Darcan S, Carr IM, Özen S, Yan Y, Hamedani MK, Gürbüz F, Mengen E, Turan İ, Ulubay A, Akkuş G, Yüksel B, Topaloglu AK, Leygue E. Idiopathic hypogonadotropic hypogonadism caused by inactivating mutations in SRA1. *J Clin Res Pediatr Endocrinol.* 2016;8:125-134. Epub 2016 Apr 18
142. Gach A, Pinkier I, Wysocka U, Sałacińska K, Salachna D, Szarras-Czapnik M, Pietrzyk A, Sakowicz A, Nykel A, Rutkowska L, Rybak-Krzyszowska M, Socha M, Jamsheer A, Jakubowski L. New findings in oligogenic inheritance of congenital hypogonadotropic hypogonadism. *Arch Med Sci.* 2022;18:353-364.
143. Neocleous V, Fanis P, Toumba M, Tanteles GA, Schiza M, Cinarli F, Nicolaidis NC, Oulas A, Spyrou GM, Mantzoros CS, Vlachakis D, Skordis N, Phylactou LA. GnRH deficient patients with congenital hypogonadotropic hypogonadism: novel genetic findings in *ANOS1*, *RNF216*, *WDR11*, *FGFR1*, *CHD7*, and *POLR3A* genes in a case series and review of the literature. *Front Endocrinol (Lausanne).* 2020;11:626.
144. Topaloglu AK, Lomniczi A, Kretzschmar D, Dissen GA, Kotan LD, McArdle CA, Koc AF, Hamel BC, Guclu M, Papatya ED, Eren E, Mengen E, Gurbuz F, Cook M, Castellano JM, Kekil MB, Mungan NO, Yuksel B, Ojeda SR. Loss-of-function mutations in PNPLA6 encoding neuropathy target esterase underlie pubertal failure and neurological deficits in Gordon Holmes syndrome. *J Clin Endocrinol Metab.* 2014;99:E2067-E2075. Epub 2014 Jul 17
145. Margolin DH, Kousi M, Chan YM, Lim ET, Schmähmann JD, Hadjivassiliou M, Hall JE, Adam I, Dwyer A, Plummer L, Aldrin SV, O'Rourke J, Kirby A, Lage K, Milunsky A, Milunsky JM, Chan J, Hedley-Whyte ET, Daly MJ, Katsanis N, Seminara SB. Ataxia, dementia, and hypogonadotropism caused by disordered ubiquitination. *N Engl J Med.* 2013;368:1992-2003. Epub 2013 May 8
146. Shi CH, Schisler JC, Rubel CE, Tan S, Song B, McDonough H, Xu L, Portbury AL, Mao CY, True C, Wang RH, Wang QZ, Sun SL, Seminara SB, Patterson C, Xu YM. Ataxia and hypogonadism caused by the loss of ubiquitin ligase activity of the U box protein CHIP. *Hum Mol Genet.* 2014;23:1013-1024. Epub 2013 Oct 9
147. Wierman ME, Xiong X, Kepa JK, Spaulding AJ, Jacobsen BM, Fang Z, Nilaver G, Ojeda SR. Repression of gonadotropin-releasing hormone promoter activity by the POU homeodomain transcription factor SCIP/Oct-6/Tst-1: a regulatory mechanism of phenotype expression? *Mol Cell Biol.* 1997;17:1652-1665.
148. Wolfe A, Kim HH, Tobet S, Stafford DE, Radovick S. Identification of a discrete promoter region of the human GnRH gene that is sufficient for directing neuron-specific expression: a role for POU homeodomain transcription factors. *Mol Endocrinol.* 2002;16:435-449.
149. Cho HJ, Gurbuz F, Stamou M, Kotan LD, Farmer SM, Can S, Tompkins MF, Mammadova J, Altincik SA, Gokce C, Catli G, Bugrul F, Bartlett K, Turan I, Balasubramanian R, Yuksel B, Seminara SB, Wray S, Topaloglu AK. *POU6F2* mutation in humans with pubertal failure alters GnRH transcript expression. *Front Endocrinol (Lausanne).* 2023;14:1203542.
150. Jee YH, Won S, Lui JC, Jennings M, Whalen P, Yue S, Temnycky AG, Barnes KM, Cheetham T, Boden MG, Radovick S, Quinton R, Leschek EW, Aguilera G, Yanovski JA, Seminara SB, Crowley WF, Delaney A, Roche KW, Baron J. DLG2 variants in patients with pubertal disorders. *Genet Med.* 2020;22:1329-1337. Epub 2020 Apr 28
151. Turan I, Demir K, Mengen E, Kotan LD, Gürbüz F, Yüksel B, Topaloglu AK. DLG2 mutations in the etiology of pubertal delay and idiopathic hypogonadotropic hypogonadism. *Horm Res Paediatr.* 2021;94:364-368. Epub 2021 Oct 25
152. Topaloglu AK, Simsek E, Kocher MA, Mammadova J, Bober E, Kotan LD, Turan I, Celiloglu C, Gurbuz F, Yuksel B, Good DJ. Inactivating NHLH2 variants cause idiopathic hypogonadotropic hypogonadism and obesity in humans. *Hum Genet.* 2022;141:295-304. Epub 2022 Jan 23
153. Good DJ, Porter FD, Mahon KA, Parlow AF, Westphal H, Kirsch IR. Hypogonadism and obesity in mice with a targeted deletion of the Nhlh2 gene. *Nat Genet.* 1997;15:397-401.
154. Alsters SI, Goldstone AP, Buxton JL, Zekavati A, Sosinsky A, Yiorakas AM, Holder S, Klaber RE, Bridges N, van Haelst MM, le Roux CW, Walley AJ, Walters RG, Mueller M, Blakemore AI. Truncating homozygous mutation of carboxypeptidase E (CPE) in a morbidly obese female with type 2 diabetes mellitus, intellectual disability and hypogonadotropic hypogonadism. *PLoS One.* 2015;10:e0131417.
155. Durmaz A, Aykut A, Atik T, Özen S, Ayyıldız Emecen D, Ata A, Işık E, Gökşen D, Çoğulu Ö, Özkinay F. A new cause of obesity syndrome associated with a mutation in the carboxypeptidase gene detected in three siblings with obesity, intellectual disability and hypogonadotropic hypogonadism. *J Clin Res Pediatr Endocrinol.* 2021;13:52-60. Epub 2020 Sep 17
156. Bosch E, Hebebrand M, Popp B, Penger T, Behring B, Cox H, Towner S, Kraus C, Wilson WG, Khan S, Krumbiegel M, Ekici AB, Uebe S, Trollmann R, Woelfle J, Reis A, Vasileiou G. BDV syndrome: an emerging syndrome with profound obesity and neurodevelopmental delay resembling Prader-Willi syndrome. *J Clin Endocrinol Metab.* 2021;106:3413-3427.
157. Bernard G, Chouery E, Putorti ML, Tétreault M, Takanoashi A, Carosso G, Clément I, Boespflug-Tanguy O, Rodriguez D, Delague V, Abou Ghoch J, Jaikh N, Dorboz I, Fribourg S, Teichmann M, Megarbane A, Schiffmann R, Vanderver A, Brais B. Mutations of POLR3A encoding a catalytic subunit of RNA polymerase Pol III cause a recessive hypomyelinating leukodystrophy. *Am J Hum Genet.* 2011;89:415-423.
158. Tétreault M, Choquet K, Orcesi S, Tonduti D, Balottin U, Teichmann M, Fribourg S, Schiffmann R, Brais B, Vanderver A, Bernard G. Recessive mutations in POLR3B, encoding the second largest subunit of Pol III, cause a rare hypomyelinating leukodystrophy. *Am J Hum Genet.* 2011;89:652-655. Epub 2011 Oct 27

159. Choquet K, Pinard M, Yang S, Moir RD, Poitras C, Dicaire MJ, Sgarioni N, Larivière R, Kleinman CL, Willis IM, Gauthier MS, Coulombe B, Brais B. The leukodystrophy mutation Polr3b R103H causes homozygote mouse embryonic lethality and impairs RNA polymerase III biogenesis. *Mol Brain*. 2019;12:59.
160. Schäfer U, Seibold S, Schneider A, Neugebauer E. Isolation and characterisation of the human rab18 gene after stimulation of endothelial cells with histamine. *FEBS Lett*. 2000;466:148-154.
161. Abdel-Hamid MS, Abdel-Ghafar SF, Ismail SR, Desouky LM, Issa MY, Effat LK, Zaki MS. Micro and Martsolf syndromes in 34 new patients: refining the phenotypic spectrum and further molecular insights. *Clin Genet*. 2020;98:445-456.
162. Handley MT, Morris-Rosendahl DJ, Brown S, Macdonald F, Hardy C, Bem D, Carpanini SM, Borck G, Martorell L, Izzi C, Faravelli F, Accorsi P, Pinelli L, Basel-Vanagaite L, Peretz G, Abdel-Salam GM, Zaki MS, Jansen A, Mowat D, Glass I, Stewart H, Mancini G, Lederer D, Roscioli T, Giuliano F, Plomp AS, Rolfs A, Graham JM, Seemanova E, Poo P, García-Cazorla A, Ederly P, Jackson IJ, Maher ER, Aligianis IA. Mutation spectrum in RAB3GAP1, RAB3GAP2, and RAB18 and genotype-phenotype correlations in warburg micro syndrome and Martsolf syndrome. *Hum Mutat*. 2013;34:686-696.
163. Xu W, Plummer L, Quinton R, Swords F, Crowley WF, Seminara SB, Balasubramanian R. Hypogonadotropic hypogonadism due to variants in RAB3GAP2: expanding the phenotypic and genotypic spectrum of Martsolf syndrome. *Cold Spring Harb Mol Case Stud*. 2020;6:a005033.
164. Bem D, Yoshimura S, Nunes-Bastos R, Bond FC, Kurian MA, Rahman F, Handley MT, Hadzhiev Y, Masood I, Straatman-Iwanowska AA, Cullinane AR, McNeill A, Pasha SS, Kirby GA, Foster K, Ahmed Z, Morton JE, Williams D, Graham JM, Dobyns WB, Burglen L, Ainsworth JR, Gissen P, Müller F, Maher ER, Barr FA, Aligianis IA. Loss-of-function mutations in RAB18 cause Warburg micro syndrome. *Am J Hum Genet* Bem D, Yoshimura S, Nunes-Bastos R, Bond FC, Kurian MA, Rahman F, Handley MT, Hadzhiev Y, Masood I, Straatman-Iwanowska AA, Cullinane AR, McNeill A, Pasha SS, Kirby GA, Foster K, Ahmed Z, Morton JE, Williams D, Graham JM, Dobyns WB, Burglen L, Ainsworth JR, Gissen P, Müller F, Maher ER, Barr FA, Aligianis IA. Loss-of-function mutations in RAB18 cause Warburg micro syndrome. *Am J Hum Genet*. 2011;88:499-507.
165. Tata B, Huijbregts L, Jacquier S, Csaba Z, Genin E, Meyer V, Leka S, Dupont J, Charles P, Chevenne D, Carel JC, Léger J, de Roux N. Haploinsufficiency of Dmxl2, encoding a synaptic protein, causes infertility associated with a loss of GnRH neurons in mouse. *PLoS Biol*. 2014;12:e1001952.
166. Lippincott MF, Xu W, Smith AA, Miao X, Lafont A, Shennib O, Farley GJ, Sabbagh R, Delaney A, Stamou M, Plummer L, Salnikov K, Georgopoulos NA, Mericq V, Quinton R, Mau-Them FT, Nambot S, Hamad A, Brittain H, Tooze RS, Calpena E, Wilkie AOM, Willems M, Crowley WF, Balasubramanian R, Lamarche-Vane N, Davis EE, Seminara SB. The p190 RhoGAPs, ARHGAP35, and ARHGAP5 are implicated in GnRH neuronal development: Evidence from patients with idiopathic hypogonadotropic hypogonadism, zebrafish, and in vitro GAP activity assay. *Genet Med*. 2022;24:2501-2515. Epub 2022 Sep 30
167. Matthews CH, Borgato S, Beck-Peccoz P, Adams M, Tone Y, Gambino G, Casagrande S, Tedeschini G, Benedetti A, Chatterjee VK. Primary amenorrhoea and infertility due to a mutation in the beta-subunit of follicle-stimulating hormone. *Nat Genet*. 1993;5:83-86.
168. Layman LC, Lee EJ, Peak DB, Namnoum AB, Vu KV, van Lingen BL, Gray MR, McDonough PG, Reindollar RH, Jameson JL. Delayed puberty and hypogonadism caused by mutations in the follicle-stimulating hormone beta-subunit gene. *N Engl J Med*. 1997;337:607-611.
169. Layman LC, Porto AL, Xie J, da Motta LA, da Motta LD, Weiser W, Sluss PM. FSH beta gene mutations in a female with partial breast development and a male sibling with normal puberty and azoospermia. *J Clin Endocrinol Metab*. 2002;87:3702-3707.
170. Sun L, Peng Y, Sharrow AC, Iqbal J, Zhang Z, Papachristou DJ, Zaidi S, Zhu LL, Yaroslavskiy BB, Zhou H, Zallone A, Sairam MR, Kumar TR, Bo W, Braun J, Cardoso-Landa L, Schaffler MB, Moonga BS, Blair HC, Zaidi M. FSH directly regulates bone mass. *Cell*. 2006;125:247-260.
171. Axelrod L, Neer RM, Kliman B. Hypogonadism in a male with immunologically active, biologically inactive luteinizing hormone: an exception to a venerable rule. *J Clin Endocrinol Metab*. 1979;48:279-287.
172. Weiss J, Axelrod L, Whitcomb RW, Harris PE, Crowley WF, Jameson JL. Hypogonadism caused by a single amino acid substitution in the beta subunit of luteinizing hormone. *N Engl J Med*. 1992;326:179-183.
173. Basciani S, Watanabe M, Mariani S, Passeri M, Persichetti A, Fiore D, Scotto d'Abusco A, Caprio M, Lenzi A, Fabbri A, Gnassi L. Hypogonadism in a patient with two novel mutations of the luteinizing hormone β -subunit gene expressed in a compound heterozygous form. *J Clin Endocrinol Metab*. 2012;97:3031-3038. Epub 2012 Jun 21
174. Yang X, Ochin H, Shu L, Liu J, Shen J, Liu J, Lin C, Cui Y. Homozygous nonsense mutation Trp28X in the LHB gene causes male hypogonadism. *J Assist Reprod Genet*. 2018;35:913-919. Epub 2018 Feb 23
175. Zhou C, Niu Y, Xu H, Li Z, Wang T, Yang W, Wang S, Wang DW, Liu J. Mutation profiles and clinical characteristics of Chinese males with isolated hypogonadotropic hypogonadism. *Fertil Steril*. 2018;110:486-495.
176. Ma X, Dong Y, Matzuk MM, Kumar TR. Targeted disruption of luteinizing hormone beta-subunit leads to hypogonadism, defects in gonadal steroidogenesis, and infertility. *Proc Natl Acad Sci U S A*. 2004;101:17294-17299. Epub 2004 Nov 29
177. Clarkson J, Han SY, Piet R, McLennan T, Kane GM, Ng J, Porteous RW, Kim JS, Colledge WH, Iremonger KJ, Herbison AE. Definition of the hypothalamic GnRH pulse generator in mice. *Proc Natl Acad Sci U S A*. 2017;114:E10216-E10223. Epub 2017 Nov 6
178. Lehman MN, Coolen LM, Goodman RL. Minireview: kisspeptin/neurokinin B/dynorphin (KNDy) cells of the arcuate nucleus: a central node in the control of gonadotropin-releasing hormone secretion. *Endocrinology*. 2010;151:3479-3489. Epub 2010 May 25
179. Uenoyama Y, Tsukamura H. KNDy neurones and GnRH/LH pulse generation: Current understanding and future aspects. *J Neuroendocrinol*. 2023;35:e13285. Epub 2023 May 26
180. Topaloglu AK, Semple RK. Neurokinin B signalling in the human reproductive axis. *Mol Cell Endocrinol*. 2011;346:57-64. Epub 2011 Jul 22
181. Navarro VM, Gottsch ML, Chavkin C, Okamura H, Clifton DK, Steiner RA. Regulation of gonadotropin-releasing hormone secretion by kisspeptin/dynorphin/neurokinin B neurons in the arcuate nucleus of the mouse. *J Neurosci*. 2009;29:11859-11866.
182. Pinilla L, Aguilar E, Dieguez C, Millar RP, Tena-Sempere M. Kisspeptins and reproduction: physiological roles and regulatory mechanisms. *Physiol Rev*. 2012;92:1235-1316.
183. Topaloglu AK, Reimann F, Guclu M, Yalin AS, Kotan LD, Porter KM, Serin A, Mungan NO, Cook JR, Imamoglu S, Akalin NS, Yuksel B, O'Rahilly S, Semple RK. TAC3 and TACR3 mutations in familial hypogonadotropic hypogonadism reveal a key role for Neurokinin B in the central control of reproduction. *Nat Genet*. 2009;41:354-358. Epub 2008 Dec 11
184. Nakamura K, Morrison SF. A thermosensory pathway mediating heat-defense responses. *Proc Natl Acad Sci U S A*. 2010;107:8848-8853. Epub 2010 Apr 26
185. Mittelman-Smith MA, Williams H, Krajewski-Hall SJ, McMullen NT, Rance NE. Role for kisspeptin/neurokinin B/dynorphin (KNDy) neurons in cutaneous vasodilatation and the estrogen modulation of body temperature. *Proc Natl Acad Sci U S A*. 2012;109:19846-19851. Epub 2012 Nov 12
186. Mittelman-Smith MA, Krajewski-Hall SJ, McMullen NT, Rance NE. Neurokinin 3 receptor-expressing neurons in the median preoptic nucleus modulate heat-dissipation effectors in the female rat. *Endocrinology*. 2015;156:2552-2562. Epub 2015 Mar 31

-
187. Fraser GL, Obermayer-Pietsch B, Laven J, Griesinger G, Pintiaux A, Timmerman D, Fauser BCJM, Lademacher C, Combalbert J, Hoveyda HR, Ramael S. Randomized controlled trial of neurokinin 3 receptor antagonist fezolinetant for treatment of polycystic ovary syndrome. *J Clin Endocrinol Metab.* 2021;106:e3519-e3532.
188. Comninos AN, Dhillon WS. Neurokinin 3 receptor antagonism for menopausal hot flashes. *Cell.* 2023;186:3332.
189. Skorupskaitė K, George JT, Veldhuis JD, Anderson RA. Neurokinin B regulates gonadotropin secretion, ovarian follicle growth, and the timing of ovulation in healthy women. *J Clin Endocrinol Metab.* 2018;103:95-104.
190. Skorupskaitė K, George JT, Veldhuis JD, Millar RP, Anderson RA. Kisspeptin and neurokinin B interactions in modulating gonadotropin secretion in women with polycystic ovary syndrome. *Hum Reprod.* 2020;35:1421-1431.
191. Ibáñez L, Ong K, Ferrer A, Amin R, Dunger D, de Zegher F. Low-dose flutamide-metformin therapy reverses insulin resistance and reduces fat mass in nonobese adolescents with ovarian hyperandrogenism. *J Clin Endocrinol Metab.* 2003;88:2600-2606.